

Datasheet for 000-001-215

Green Fluorescent Protein (GFP) Control Protein

Overview

Description:	Recombinant Green Fluorescent Protein (GFP) Control Protein - 000-001-215
Item No.:	000-001-215
Size:	100 µg
Applications:	SDS-PAGE, WB, Microarray
Origin:	Aequorea victoria
Expressed in:	E. coli

Product Details

Background:	Green Fluorescent Protein (GFP) is used as a positive control when detecting GFP fusion proteins in Anti-GFP antibody assays. Rockland offers multiple purified hosts and conjugated Anti-GFP that can detect GFP by ELISA (sandwich or capture) or western blot. Biotin conjugated polyclonal Anti-GFP used in a sandwich ELISA is well suited to titrate GFP in solution when used in combination with monoclonal Anti-GFP (p/n 600-301-215), using either form of the antibody as the capture or detection antibody.
Synonyms:	control protein, rGFP protein, GFP fusion protein, GFP control
Species of Origin:	Aequorea victoria
Expressed in:	E. coli

Target Details

Purity/Specificity:	GFP control protein was synthesized with an amino terminal 6XHIS tag and was expressed in E. coli and purified by sequential hydrophobic interaction chromatography, DEAE chromatography and size-exclusion chromatography. Assay by immunoelectrophoresis resulted in a single precipitin arc against Anti-GFP. Analysis by SDS-PAGE resulted in a pattern consistent with purified GFP and was estimated to be greater than 90% pure.
Relevant Links:	UniProtKB - P42212

Application Details

Tested Applications:	SDS-PAGE, WB
Suggested Applications:	Microarray (Based on references)
Application Note:	GFP Control is suitable as a control for polyclonal or monoclonal Anti-GFP in immunological assays. Researchers should determine optimal titers for applications. The molecular weight of GFP is 28 kDa. Green Fluorescent Protein has been tested in SDS-page, western blot, and immunoprecipitation.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	User Optimized
IP:	User Optimized
WB:	User Optimized


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by BCA assay
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling


Shipping Condition:	Dry Ice
Storage Condition:	Store GFP at -20 °C prior to opening. Aliquot contents and freeze at -20 °C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4 °C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is six (6) months from date of receipt.

Images




Western Blot

Western Blot of anti-GFP monoclonal antibody. Lane 1: 64pg of recombinant GFP protein (p/n 000-001-215) were spiked into a HeLa cell-derived lysates (p/n W09-000-364). Lane 2: 32pg of recombinant GFP protein were spiked into a HeLa cell-derived lysates. Lane 3: 16pg of recombinant GFP protein were spiked into a HeLa cell-derived lysates. Lane 4: 8pg of recombinant GFP protein were spiked into a HeLa cell-derived lysates. Lane 5: 4pg of recombinant GFP protein were spiked into a HeLa cell-derived lysates. Lane 6: 2pg of recombinant GFP protein were spiked into a HeLa cell-derived lysates. Lane 7: 1pg of recombinant GFP protein were spiked into a HeLa cell-derived lysates. Lane 8: 0pg of recombinant GFP protein were spiked into a HeLa cell-derived lysates. Primary antibody: anti-GFP monoclonal antibody at 1:400 for overnight at 4°C. Secondary antibody: HRP-conjugated anti-Mouse IgG (p/n 610-4302) was performed at a dilution of 1:20,000 for 1h at 4°C. Block: TTBS (p/n MB-013) supplemented with 1% BSA (p/n BSA-50) for 1 h at 4°C. Predicted/Observed size: 27 kDa for GFP. Other band(s): none.



SDS-PAGE

SDS-PAGE results of Recombinant GFP Control Protein. Lane 1: Recombinant GFP Control Protein Reduced (1.0µg). Lane 2: Opal Prestained Molecular Weight Marker (p/n MB-210-0500). Lane 3: Recombinant GFP Control Protein Non-Reduced (1.0µg). Predicted MW: ~28kDa. Observed MW: ~28kDa doublet, ~60kDa dimer. 4-20% Gel Coomassie Stained.



Western Blot

Immunoprecipitation/Western Blot using GFP Protein. Lane 1: Opal Prestained Molecular Weight Marker (p/n MB-210-0500). Lane 2: GFP Input (p/n 000-001-215) Reduced [10 μ L]. Primary IP Antibody: Mouse Anti-GFP (p/n 600-301-215) at 10 μ g overnight at 2-8°C. Secondary Antibody: TrueBlot Anti-Mouse Ig IP Agarose Beads (p/n 00-8811-25) at 500 μ g for 1hr at RT. Buffer: BlockOut Buffer (p/n MB-073) for 30 mins at RT. Exposure: 7 sec.

References

- Dolan ME et al. First site-specific conjugation method for native goat IgG antibodies via glycan remodeling at the conserved Fc region. *Antib Ther.* (2024)
- Dufour YS et al. Direct correlation between motile behavior and protein abundance in single cells. *PLoS Comput Biol.* (2016)
- Pla-Roca, M et al. Antibody colocalization microarray: a scalable technology for multiplex protein analysis in complex samples. *Molecular & Cellular Proteomics : MCP* (2012)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.