

Datasheet for 001-0305

## Bovine IgG Fab Peroxidase

### Overview

|                     |  |
|---------------------|--|
| <b>Description:</b> | Bovine IgG Fab Fragment Peroxidase Conjugated - 001-0305 |
| <b>Item No.:</b>    | 001-0305   |
| <b>Size:</b>        | 1 mg   |
| <b>Origin:</b>      | Bovine   |

### Product Details

|                           |   |
|---------------------------|---|
| <b>Background:</b>        | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the compliment cascade, and opsonization for phagocytosis. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. |
| <b>Synonyms:</b>          | Bovine IgG Fab peroxidase conjugation, Bovine IgG Fab HRP conjugation, peroxidase conjugated Bovine IgG Fab, HRP conjugated Bovine IgG Fab fragment   |
| <b>Species of Origin:</b> | Bovine  |
| <b>Conjugate:</b>         | Peroxidase (HRP)  |
| <b>Format:</b>            | IgG Fab   |
| <b>Type:</b>              | Native Protein  |

### Target Details

|                            |  |
|----------------------------|--|
| <b>Purity/Specificity:</b> | This product was prepared from normal serum by delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti- Bovine IgG, anti-Bovine IgG F(ab') <sub>2</sub> and anti-Bovine Serum. No reaction was observed against anti-Bovine IgG F(c) or anti-Papain. |
|----------------------------|--|

### Application Details

|                          |   |
|--------------------------|---|
| <b>Application Note:</b> | Bovine IgG F(ab) fragment peroxidase conjugation can be utilized as a control or standard |
|--------------------------|---|

reagent in Western Blotting and ELISA experiments.

---

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

---

## Formulation

---

**Physical State:** Lyophilized

---

**Concentration:** 1.0 mg/mL by UV absorbance at 280 nm

---

**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

---

**Preservative:** 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

---

**Stabilizer:** 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

---

**Reconstitution Volume:** 1.0 mL

---

**Reconstitution Buffer:** Restore with deionized water (or equivalent)

---

## Shipping & Handling

---

**Shipping Condition:** Ambient

---

**Storage Condition:** Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

---

**Expiration:** Expiration date is one (1) year from date of receipt.

---

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

