Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 005-0051

Goat IgG F(ab')2 Agarose

Overview

Description:	Goat IgG F(ab')2 Fragment Agarose Conjugated - 005-0051
Item No.:	005-0051
Size:	5 mg
Applications:	SDS-PAGE
Origin:	Goat

Product Details

Background:	Goat IgG F(ab')2 Fragment Agarose Conjugated is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlled conditions of temperature, time and pH. Goat IgG F(ab')2 molecules lack the Fc portion of Goat IgG and therefore receptors that bind Goat IgG Fc will not bind Goat IgG F(ab')2 molecules. This product possesses the F(ab')2 fragment, recognized by the two F(ab) fragments yielded from the digestion of the antibody below the disulfide bond hinge region followed by agarose conjugation.
Synonyms:	Goat IgG F(ab')2 fragment Agarose Conjugate, Goat IgG Fab2 Agarose
Species of Origin:	Goat
Conjugate:	Agarose
Format:	IgG F(ab')2
Type:	Native Protein

Target Details

Purity/Specificity:	This product is an purified Goat IgG isolated from normal Goat Serum digested by the enzyme pepsin and purified by chromatography coupled to activated agarose. Sufficient antibody capacity is provided to bind a minimum of 5.0 mg of anti-Goat IgG.

Application Details

Tested Applications:	SDS-PAGE

Application Note:	Goat IgG F(ab')2 Agarose has been tested by SDS-PAGE and is suitable for immunoprecipitation and Immunoaffinity purifications.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
IP:	User Optimized


Formulation

Physical State:	Suspension of agarose beads
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. DO NOT FREEZE.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



SDS-PAGE
SDS-PAGE of Goat IgG Whole Molecule Rhodamine Conjugated (p/n 005-0002). Lane M: 5 µL Opal Prestained Marker (p/n MB-210-0500). Lane 1: Reduced Goat IgG Whole Molecule Rhodamine Conjugated (p/n 005-0002). Lane 2: Reduced Goat IgG F(c) Fragment (p/n 005-0103). Lane 3: Reduced Goat IgG F(ab) Fragment (p/n 005-0105). Lane 4: Reduced Goat IgM Whole Molecule (p/n 005-0107). Load: 1 µg for IgG, F(c) and F(ab); 3 µg for IgM. Predicted/Observed size: IgG at 50 and 25 kDa; F(c) at 25 kDa; F(ab) at 25 kDa; IgM at 70 and 23 kDa. Observed F(c) Fragment migrates slightly higher.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.