

**Datasheet for 008-0204****Horse IgG F(ab')<sub>2</sub> Fluorescein****Overview**

|                     |                                                                          |
|---------------------|--------------------------------------------------------------------------|
| <b>Description:</b> | Horse IgG F(ab') <sub>2</sub> Fragment Fluorescein Conjugated - 008-0204 |
| <b>Item No.:</b>    | 008-0204                                                                 |
| <b>Size:</b>        | 1 mg                                                                     |
| <b>Origin:</b>      | Horse                                                                    |

**Product Details**

|                           |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|---------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Background:</b>        | Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well as fungi and facilitates their destruction or neutralization via agglutination (and thereby immobilizing them), activation of the complement cascade, and opsonization for phagocytosis. This product possesses the F(ab') <sub>2</sub> fragment, recognized by the two F(ab) fragments yielded from the digestion of the antibody below the disulfide bond hinge region. |
| <b>Synonyms:</b>          | Horse IgG F(ab') <sub>2</sub> fragment fluorescein conjugated, Horse IgG F(ab') <sub>2</sub> fragment FITC conjugated                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Species of Origin:</b> | Horse                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Conjugate:</b>         | Fluorescein (FITC)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Format:</b>            | IgG F(ab') <sub>2</sub>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| <b>Type:</b>              | Native Protein                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>F/P Ratio:</b>         | 2.3                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |

**Target Details**

|                            |                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Purity/Specificity:</b> | This product was prepared from normal serum by delipidation, salt fractionation and ion change chromatography followed by pepsin digestion and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Horse IgG, anti-Horse IgG F(ab') <sub>2</sub> and anti-Horse Serum. No reaction was observed against anti-Horse IgG F(c) or anti-Pepsin. |
|----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

**Application Details**

|                          |                                                                                                                                                                                                                                                                                                         |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Application Note:</b> | Horse IgG F(ab') <sub>2</sub> fragment Fluorescein is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. |
| <b>Assay Dilutions:</b>  | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.                                                                                                                                                                                                         |

## Formulation

|                               |                                                                        |
|-------------------------------|------------------------------------------------------------------------|
| <b>Physical State:</b>        | Lyophilized                                                            |
| <b>Concentration:</b>         | 1.0 mg/mL by UV absorbance at 280 nm                                   |
| <b>Buffer:</b>                | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2                |
| <b>Preservative:</b>          | 0.01% (w/v) Thimerosal                                                 |
| <b>Stabilizer:</b>            | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| <b>Reconstitution Volume:</b> | 1.0 mL                                                                 |
| <b>Reconstitution Buffer:</b> | Restore with deionized water (or equivalent)                           |

## Shipping & Handling

|                            |                                                                                                                                                                                                                                                                                                                                                 |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Shipping Condition:</b> | Ambient                                                                                                                                                                                                                                                                                                                                         |
| <b>Storage Condition:</b>  | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| <b>Expiration:</b>         | Expiration date is one (1) year from date of receipt.                                                                                                                                                                                                                                                                                           |

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

