

Datasheet for 009-001-C20-0100**rHuman RANTES (CCL5) Protein****Overview**

Description:	Human RANTES (CCL5) Recombinant Protein - 009-001-C20-0100
Item No.:	009-001-C20-0100
Size:	100 µg
Applications:	Cellular Assay
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	Regulated Upon Activation Normal T cell Express Sequence (RANTES), also called CCL5, is a chemokine produced by T cells three to five days after activation. RANTES is a promiscuous chemokine that signals through several G protein-coupled receptors, CCR5, CCR3, CCR1 and US28 (a viral receptor encoded by human CMV). The main function of RANTES is to recruit immune cells to the site of inflammation. Recombinant human RANTES is a non-glycosylated protein, containing 68 amino acids, with a molecular weight of 7.8 kDa.
Synonyms:	EoCP, Eosinophil chemotactic cytokine, SIS-delta, Small-inducible cytokine A5, T cell-specific protein P228 (TCP228), T-cell-specific protein RANTES
Species of Origin:	Human
Expressed in:	E. coli
Type:	Recombinant Protein
Low Endotoxin:	Yes

Target Details

Gene Name:	CCL5
Purity/Specificity:	RANTES (CCL5) purity was determined to be greater than 98% as determined by HPLC, analysis by UV-Spectroscopy at 280nm, and by reducing and non-reducing SDS-PAGE.
Relevant Links:	• UniProtKB - P13501

Application Details

Suggested Applications:	Cellular Assay (Based on references)
Application Note:	RANTES Recombinant Protein has been tested by biologic activity and is suitable as a control for polyclonal or monoclonal anti-RANTES in immunological assays.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	Endotoxin Level: Measured by kinetic LAL analysis and is typically ≤ 1 EU/ μ g protein. Biologic Activity: The activity is determined by the ability to chemoattract human monocytes, neutrophils, THP-1 or primary T cells at 1-8 ng/mL.


Formulation

Physical State:	Lyophilized
Buffer:	0.1% Trifluoroacetic acid
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	100 μ L
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
Expiration:	Expiration date is six (6) months from date of receipt.

Images



Figure

Bioactivity of Human RANTES (CCL5) Recombinant Protein. Human THP-1 cells were allowed to migrate to Human RANTES at (0, 0.1, 1, 10, 100 and 1000 ng/mL). After 45 minutes, cells that migrated were counted using a luminescent substrate and displayed on the bar graph above. Significant increases in migration over basal levels were seen in response to Human RANTES detectable starting at 10 ng/mL. These results are similar to results expected from primary human monocytes.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.