

Datasheet for 009-001-H86S**IKK gamma protein-GST fusion****Overview**

Description:	IKK gamma recombinant protein-GST fusion protein - 009-001-H86S
Item No.:	009-001-H86S
Size:	20 µg
Origin:	Human
Expressed in:	Sf9 cells

Product Details

Background:	IKKgamma is a serine/threonine protein kinase that phosphorylates the I-kappa-B protein which is an inhibitor of the transcription factor NF-kappa-B complex. Phosphorylation of I-kappa-B protein triggers the degradation of the inhibitor via the ubiquitination pathway, thereby activating NF-kappa-B complex. IKKgamma forms dimers and trimers and interacts preferentially with IKKbeta but not IKKalpha (1). IKKgamma associates with activated ATM after the induction of DNA double-strand breaks (2). ATM phosphorylates serine-85 of IKKgamma to promote its ubiquitin-dependent nuclear export. IKKgamma Protein is ideal for investigators involved in Signaling Proteins, Cellular Proteins, Cancer, Inflammation, Neurobiology, Ser/Thr Kinases, and WNT Signaling research.
Synonyms:	IKBKG, AMCBX1, FIP-3, FIP3, Fip3p, IKK-gamma, IP, IP1, IP2, IPD2, NEMO, NF-kappa-B essential modulator, NEMO, FIP-3, Ikb kinase-associated protein 1, IKKAP1, Inhibitor of nuclear factor kappa-B kinase subunit gamma, I-kappa-B kinase subunit gamma, IKK-gamma, IKKG, Ikb kinase subunit gamma, NF-kappa-B essential modifier
Species of Origin:	Human
Expressed in:	Sf9 cells
Type:	Recombinant Protein

Target Details

Gene Name:	IKBKG
Purity/Specificity:	Recombinant full-length human IKK gamma was expressed by baculovirus in Sf9 insect cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >95% by densitometry.

Relevant Links:

- [NCBI - NM_003639](#)

Application Details

Application Note: IKK gamma Protein is stored in 50mM Tris-HCl, pH 7.5, 50mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol. IKK gamma Protein is suitable for use in Western Blot. Expect a band approximately ~73kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

WB: User Optimized

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 0.2 µg/µL

Buffer: See application note.

Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

Expiration: Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

