Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 009-001-S19S

I kappaB beta protein-GST fusion

Overview

Description:	I kappaB beta recombinant protein-GST fusion protein - 009-001-S19S
Item No.:	009-001-S19S
Size:	20 µg
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	IkB-beta is part of the NFkB complex and it inactivates NFkB complex by binding to it and trapping it in the cytoplasm. Phosphorylation of serine residues on IkBβ mediated by IkB kinases leads to its destruction via the ubiquitination pathway thereby allowing activation of the NFkB complex which translocates into the nucleus and binds DNA at kappa-B-binding motifs (1). The PEST domain of IkBβ can interact with two proteins KBRAS1 and KBRAS2 and this interaction can decrease the rate of degradation IkBβ. IkBβ has been shown to participate in multiple signaling pathways including adipocytokine signaling pathway, B-cell receptor signaling pathway, T-cell receptor signaling pathway and Hypoxia (2). IkBβ Protein is ideal for investigators involved in Signaling Proteins, Transcription Proteins, AKT/PKB Pathway, Apoptosis/Autophagy, Inflammation, and p38 Pathway research.
Synonyms:	IKBB, NFKBIB, TRIP9, NF-kappa-B inhibitor beta , NF-kappa-BIB, I-kappa-B-beta, IkB-B, IkB-beta, IkappaBbeta, Thyroid receptor-interacting protein 9, TR-interacting protein 9, TRIP-9
Species of Origin:	Human
Expressed in:	E. coli
Type:	Recombinant Protein

Target Details

Gene Name:	NFKBIB
Purity/Specificity:	Recombinant full-length human IkB-beta was expressed in E. coli cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >90% by densitometry.
Relevant Links:	• NCBI - NM_002503

Application Details

Application Note:	IkB-beta Protein is stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol. IkB-beta Protein is suitable for use in Western Blot and Kinase Assay. Expect a band approximately ~46kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
WB:	User Optimized

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.2 µg/µL
Buffer:	See application note.

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store product at –70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.
Expiration:	Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com
tech@rockland.com
+1 484.791.3823