

Datasheet for 009-001-S51S

Mortalin protein-GST fusion**Overview**

| | |
|----------------------|--|
| Description: | Mortalin recombinant protein-GST fusion protein - 009-001-S51S |
| Item No.: | 009-001-S51S |
| Size: | 20 µg |
| Origin: | Human |
| Expressed in: | Sf9 cells |

Product Details

| | |
|---------------------------|--|
| Background: | Mortalin is a mitochondrial chaperone and a member of the heat shock protein 70 family that is constitutively expressed in cells. Mortalin plays a central role in mitochondrial biogenesis through its capacity to direct the import of nuclear-encoded proteins into the mitochondria. Mortalin plays a role in the control of cell proliferation and elevated levels of mortalin have correlated with malignant transformation and poor cancer prognosis. Mortalin can support cancer cell resistance to complement-dependent cytotoxicity (1). Mortalin expression was decreased in the mitochondrial fraction of neurons from the substantia nigra of Parkinson disease patients (2). Mortalin Protein is ideal for investigators involved in Signaling Proteins, Cell Stress & Chaperone Proteins, Apoptosis/Autophagy, and Cancer research. |
| Synonyms: | HSPA9, CSA, GRP75, HSPA9B, MGC4500, MOT, mot-2, MOT2, MTHSP75, PBP74, Stress-70 protein, mitochondrial 75 kDa glucose-regulated protein, GRP-75, Heat shock 70 kDa protein 9, Mortalin, MOT, Peptide-binding protein 74, PBP74 |
| Species of Origin: | Human |
| Expressed in: | Sf9 cells |
| Type: | Recombinant Protein |

Target Details

| | |
|----------------------------|---|
| Gene Name: | HSPA9 |
| Purity/Specificity: | Recombinant human Mortalin (47-end) protein was expressed in Sf9 insect cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >80% by densitometry. |
| Relevant Links: | <ul style="list-style-type: none">• NCBI - NM_004134 |

Application Details

| | |
|--------------------------|---|
| Application Note: | Mortalin Protein is stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol. Mortalin Protein is suitable for use in Western Blot. Expect a band approximately ~97kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user. |
| Assay Dilutions: | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| WB: | User Optimized |

Formulation

| | |
|------------------------|-------------------------------------|
| Physical State: | Liquid (sterile filtered) |
| Concentration: | 0.2mg/mL by UV absorbance at 280 nm |
| Buffer: | See application note. |

Shipping & Handling

| | |
|----------------------------|---|
| Shipping Condition: | Dry Ice |
| Storage Condition: | Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles. |
| Expiration: | Expiration date is one (1) year from date of receipt. |

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

