

Datasheet for 009-001-S66S
p53 protein-GST fusion**Overview**

Description:	p53 recombinant protein-GST fusion protein - 009-001-S66S
Item No.:	009-001-S66S
Size:	20 µg
Origin:	Human
Expressed in:	HighFive™ cells

Product Details

Background:	p53 is a key regulator of cell growth and acts as a tumor suppressor gene. Wild-type p53 gene can suppress transformation of rat embryo fibroblasts in cell culture by other oncogenes such as the adenovirus E1A and Ras (1). DNA tumor viruses such as SV40 large T antigen and the adenovirus E1A plus E1B-55Kd proteins bind to p53 and inactivate its tumor suppressor activities leading to cellular transformation. Mutational inactivation of the p53 gene is detected in more than 50% of human cancers. Mutation of p53 renders cancer cells more resistant to current cancer therapies due to lack of p53-mediated apoptosis (2). p53 Protein is ideal for investigators involved in Signaling Proteins, Cell Cycle Proteins, Apoptosis/Autophagy, Cancer, Cardiovascular Disease, Cell Cycle, Cellular Stress, Inflammation, JNK/SAPK Pathway, Metabolic Disorder, Neurobiology, and p38 Pathway research.
Synonyms:	TP53, LFS1, TRP53, Cellular tumor antigen p53, Antigen NY-CO-13, Phosphoprotein p53, Tumor suppressor p53
Species of Origin:	Human
Expressed in:	HighFive™ cells
Type:	Recombinant Protein

Target Details

Gene Name:	TP53
Purity/Specificity:	Recombinant full-length human p53 was expressed by baculovirus in Hi5 cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >95% by densitometry.
Relevant Links:	<ul style="list-style-type: none">• NCBI - NM_000546

Application Details

Application Note:	p53 Protein is stored in 50mM Tris-HCl, pH 7.5, 50mM NaCl, 0.25mM DTT, 0.1mM EDTA, 0.1mM PMSF, 10mM glutathione, and 25% glycerol. p53 Protein is suitable for use in Western Blot and Kinase Assay. Expect a band approximately ~79kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
WB:	User Optimized

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.2 µg/µL
Buffer:	See application note.

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.
Expiration:	Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

www.rockland.com
tech@rockland.com
+1 484.791.3823