

Datasheet for 009-001-T31S**SIRT1 protein-GST fusion****Overview**

Description:	SIRT1 recombinant protein-GST fusion protein - 009-001-T31S
Item No.:	009-001-T31S
Size:	20 µg
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	SIRT1 is a member of the sirtuin family of proteins which are homologs to the yeast Sir2 protein. Sirtuin family contain a sirtuin core domain and are grouped into four classes with SIRT1 being a member of class I. SIRT1 is a stress-response and chromatin-silencing factor (1). It is an NAD(+)-dependent histone deacetylase involved in various nuclear events such as transcription, DNA replication, and DNA repair. SIRT1 protein binds and deacetylates the p53 protein (2). Expression of wild type SIRT1 in human cells reduces the transcriptional activity of p53 indicating that SIRT1 is involved in the regulation of p53 function via deacetylation. SIRT1 Protein is ideal for investigators involved in Signaling Proteins, Deacetylase/Demethylase Proteins, Cancer, Cell Cycle, and Inflammation research.
Synonyms:	SIR2L1, Sirtuin 1, NAD-dependent protein deacetylase sirtuin-1, Regulatory protein SIR2 homolog 1, SIR2-like protein 1, SirtT1 75 kDa fragment, 75SirT1
Species of Origin:	Human
Expressed in:	E. coli
Type:	Recombinant Protein

Target Details

Gene Name:	SIRT1
Purity/Specificity:	Recombinant human SIRT1 (193-end) was expressed in E. coli cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >80% by densitometry.
Relevant Links:	• NCBI - NM_012238

Application Details

Application Note:	SIRT1 Protein is stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol. SIRT1 Protein is suitable for use in Western Blot. Expect a band approximately ~105kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
WB:	User Optimized

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.2 µg/µL
Buffer:	See application note.

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.
Expiration:	Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

