

Datasheet for 009-001-T39S**SMAD3 protein-GST fusion****Overview**

Description:	SMAD3 recombinant protein-GST fusion protein - 009-001-T39S
Item No.:	009-001-T39S
Size:	20 µg
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	SMAD3 is a direct mediator of transcriptional activation by the TGFβ receptor. The activity of SMAD3 is regulated by the TGFβ receptors, and SMAD3 is phosphorylated and associated with the ligand-bound receptor complex. TGFβ stimulation leads to phosphorylation and activation of SMAD3, which form a complex with SMAD4 that accumulate in the nucleus and regulate transcription of target genes such as CDK inhibitor (1). SMAD3 containing a C-terminal truncation acts as a dominant-negative inhibitor of the normal TGFβ response. SMAD3 is a major physiologic substrate of the G1 cyclin-dependent kinases CDK4 and CDK2 (2). SMAD3 Protein is ideal for investigators involved in Signaling Proteins, Transcription Proteins, AKT/PKB Pathway, Angiogenesis, Cancer, Cell Cycle, Cellular Stress, ERK/MAPK Pathway, Inflammation, JAK/STAT Pathway, JNK/SAPK Pathway, NfκB Pathway, and WNT Signaling research.
Synonyms:	MADH3, JV15-2, HSPC193, HsT17436, MGC60396, DKFZP586N0721, DKFZp686J10186, Mothers against decapentaplegic homolog 3, MAD homolog 3, Mad3, Mothers against DPP homolog 3, hMAD-3, JV15-2
Species of Origin:	Human
Expressed in:	E. coli
Type:	Recombinant Protein

Target Details

Gene Name:	SMAD3
Purity/Specificity:	Recombinant full-length human SMAD3 was expressed in E. coli cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >90% by densitometry.

Relevant Links:

- [NCBI - NM_005902](#)

Application Details

Application Note: SMAD3 Protein is stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol. SMAD3 Protein is suitable for use in Western Blot and Kinase Assay. Expect a band approximately ~77kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

WB: User Optimized

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 0.2 µg/µL

Buffer: See application note.

Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

Expiration: Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

