

Datasheet for 009-001-T52S

STAT2 protein-GST fusion**Overview**

Description:	STAT2 recombinant protein-GST fusion protein - 009-001-T52S
Item No.:	009-001-T52S
Size:	20 µg
Origin:	Human
Expressed in:	Sf9 cells

Product Details

Background:	STAT2 (Signal transducer and activator of transcription) is a family member of STAT transcription factors that are implicated in programming gene expression in biological events as diverse as embryonic development, programmed cell death, organogenesis, innate immunity, adaptive immunity and cell growth regulation (1). JAKs can phosphorylate and activate STATs. Activation of the JAK/STAT pathway in response to cytokine action can lead to regulation of gene expression during proliferation and differentiation of the immune cells. STATs have been shown to be involved in inflammation, autoimmune and allergic diseases (2). STAT2 Protein is ideal for investigators involved in Signaling Proteins, Transcription Proteins, Apoptosis/Autophagy, Cancer, ERK/MAPK Pathway, Inflammation, and JAK/STAT Pathway research.
Synonyms:	STAT113, ISGF-3, MGC59816, P113, Signal transducer and activator of transcription 2
Species of Origin:	Human
Expressed in:	Sf9 cells
Type:	Recombinant Protein

Target Details

Gene Name:	STAT2
Purity/Specificity:	Recombinant full-length human STAT2 was expressed by baculovirus in Sf9 insect cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >90% by densitometry.
Relevant Links:	• NCBI - NM_005419

Application Details

Application Note:	STAT2 Protein is stored in 50mM Tris-HCl, pH 7.5, 150mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol. STAT2 Protein is suitable for use in Western Blot and Kinase Assay. Expect a band approximately ~138kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
WB:	User Optimized

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.2 µg/µL
Buffer:	See application note.

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.
Expiration:	Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

