

Datasheet for 009-001-W25-1000**rHuman TARC /CCL17 Protein****Overview**

Description:	Human TARC (CCL17) Recombinant Protein - 009-001-W25-1000
Item No.:	009-001-W25-1000
Size:	1 mg
Origin:	Human
Expressed in:	E. coli

Product Details

Background:	Thymus and activation regulated chemokine (TARC), also known as CCL17, is a chemokine produced by thymus tissue constitutively and activated PBMCs (mainly DCs). TARC signals through the CCR4 receptor to induce chemotaxis of Th2 cells. TARC is thought to be important in asthma and allergic diseases, along with bacterial and viral infections. Recombinant human TARC is a non-glycosylated protein, containing 71 amino acids, with a molecular weight of 8.1 kDa.
Synonyms:	Thymus and activation regulated chemokine, ABCD-2, SCYA17, CC chemokine TARC, Small-inducible cytokine A17
Species of Origin:	Human
Expressed in:	E. coli
Type:	Recombinant Protein
Low Endotoxin:	Yes

Target Details

Gene Name:	CCL17
Purity/Specificity:	TARC (CCL17) purity was determined to be greater than 98% as determined by HPLC, analysis by UV-Spectroscopy at 280nm, and by reducing and non-reducing SDS-pAGE.
Relevant Links:	• UniProtKB - Q92583

Application Details

Application Note:	TARC Recombinant Protein is suitable as a control for polyclonal or monoclonal anti-TARC in immunological assays.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	Endotoxin Level: Measured by kinetic LAL analysis and is typically ≤ 1 EU/ μ g protein. Biologic Activity: The activity is determined by the ability to chemoattract human T cells at 2-40 ng/mL.

Formulation

Physical State:	Lyophilized
Buffer:	0.1% Trifluoroacetic acid
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
Expiration:	Expiration date is six (6) months from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.