

Datasheet for 009-0034**Human Transferrin Rhodamine****Overview**

Description:	Human Transferrin Rhodamine Conjugated - 009-0034
Item No.:	009-0034
Size:	1 mg
Applications:	Dot Blot, IF, Multiplex
Origin:	Human

Product Details

Background:	Human transferrin is encoded by the TF gene and is an iron-binding blood plasma glycoprotein that controls the level of free iron in biological fluids. Human transferrin binds iron very tightly but reversibly. Human transferrin is the most important iron pool in mammals. Human transferrin has a molecular weight of around 80 kDa and contains 2 specific high-affinity Fe(III) binding sites. The affinity of Human transferrin for Fe(III) is extremely high but decreases progressively with decreasing pH below neutrality.
Synonyms:	Human transferrin Rhodamine conjugation, TRITC conjugated transferrin
Species of Origin:	Human
Conjugate:	Rhodamine (TRITC)
Format:	Transferrin
Type:	Native Protein
F/P Ratio:	2.27

Target Details

Purity/Specificity:	This product was prepared from normal serum by delipidation, salt fractionation, selective precipitation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Human Transferrin and anti-Human Serum.
Relevant Links:	• GenelD - 7018 • UniProtKB - Q06AH7

- [NCBI - AAB22049.1](#)

Application Details

Tested Applications:	Dot Blot
Suggested Applications:	IF, Multiplex (Based on references)
Application Note:	Human transferrin rhodamine conjugation has been tested by dot blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA), fluorescent western blotting, multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.


Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/ml Polyethylene Glycol (PEG-8000)
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling


Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images




Immunofluorescence Microscopy

COMMD1–PtdIns(4,5)P₂ interaction measured by fluorescence quenching of all aromatics. (C) Example images showing colocalization of ATP7B with wild type, T174M and K167/173E COMMD1. ATP7B is in green and COMMD1 variants in magenta as indicated. Fig. 6.. PMID: 31515276.




Immunofluorescence Microscopy

Colocalization of ATP7B with TGN46 or LAMP1. Cells were left untreated (basal medium), treated with 10 μM TTM (low copper) or treated with 10, 100 or 200 μM CuCl₂. (A,B) Merged images show ATP7B in green, TGN46 (A) or LAMP1 (B) in magenta and the nucleus in blue; pixel overlap is shown in white. Fig. 1. PMID: 31515276.



Immunofluorescence Microscopy

Rab5c over activation leads to excessive endolysosomal trafficking of Notch ligands and receptor. (A) TRITC-conjugated TF internalization assay in Hela cells transfected with control empty pCS2 or pCS2-rab5c CA plasmid. Representative pictures were shown. Scale bar, 10 μm. (B) Quantitative fluorescence intensity of intracellular TRITC-TF in control empty pCS2 and pCS2-rab5c CA transfected cells, n = 8 cells for each group. Error bars, mean ± SD. P value was calculated by Student t test, ***P < 0.001. CA, constitutively active; TF, transferrin; TRITC, tetramethylrhodamine. S6 Fig. PMID: 32275659.




Immunofluorescence Microscopy

Rab5c function is in an EC autonomous manner. (A) TRITC-conjugated TF internalization assay in HeLa cells transfected with empty pCS2 or pCS2-rab5c DN plasmids. Representative pictures were shown. Scale bar, 10 μ m. (B) Quantitative fluorescence intensity of intracellular TRITC-TF in empty pCS2 or pCS2-rab5c DN transfected HeLa cells, n = 8 cells for each group. Error bars, mean $\pm$ SD. P value was calculated by Student t test, ***P < 0.001. DN, dominant-negative; TF, transferrin; TRITC, tetramethylrhodamine. S3 Fig. PMID: 32275659.

Dot Blot

Dot Blot of Rhodamine Conjugated Human Transferrin. Dotted directly with Rhodamine Conjugated Human Transferrin at following concentrations. Load: Lane 1 - 50ng Lane 2 - 16.67ng Lane 3 - 5.56ng Lane 4 - 1.85ng Lane 5 - 0.62ng Primary antibody: none Secondary antibody: none Block: MB-070 for 1 HR at RT.



References

- Bheri S et al. Composition of cardiac-derived extracellular vesicles changes with vesicle origin and determines uptake. *Front Cardiovasc Med.* (2025)
- Heng J et al. Rab5c-mediated endocytic trafficking regulates hematopoietic stem and progenitor cell development via Notch and AKT signaling. *PLoS Biol.* (2020)
- Stewart DJ, Short KK, Maniaci BN, Burkhead JL. COMMD1 and PtdIns(4,5)P2 interaction maintain ATP7B copper transporter trafficking fidelity in HepG2 cells. *J Cell Sci.* (2019)

Disclaimer

No test method can provide total assurance that the hepatitis B virus, hepatitis C virus, human immunodeficiency virus, or any other infectious agents are absent. Thus, all blood products, including purified proteins derived from human blood sources, should be handled at Biosafety Level 2 as recommended by the CDC\NIH manual entitled Biosafety in Microbiological and Biomedical Laboratories for potentially infectious human serum, blood specimens or proteins derived from same. Source material for the human blood product supplied to your facility has been tested for the detection of HIV antibody, Hepatitis B surface antigen, antibody to Hepatitis C, HIV 1 antigen(s), antibody to HTLV - I/II, and syphilis by FDA guidelines. All units were found to be non-reactive/negative for these tests. All human blood source material is collected in FDA licensed centers and is tested with FDA approved test kits.

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.