

**Datasheet for 010-001-C55-0100**  
**rMouse EBI-3 Protein****Overview**

<b>Description:</b>	Mouse Epstein-Barr Virus Induced Gene 3 Recombinant Protein - 010-001-C55-0100
<b>Item No.:</b>	010-001-C55-0100
<b>Size:</b>	100 µg
<b>Applications:</b>	SDS-PAGE
<b>Origin:</b>	Mouse
<b>Expressed in:</b>	E. coli

**Product Details**

<b>Background:</b>	Epstein-Barr Virus Induced Gene-3 (EBI3), is a secreted glycoprotein belonging to the hematopoietin receptor family and is related to the p40 subunit of IL-12. EBI3 was identified by its induced expression in B-lymphocytes in response to Epstein-Barr virus infection. EBI3 forms heterodimers with p28 to form IL-27 and with p35 to form IL-35. Both IL-27 and IL-35 have anti-inflammatory and regulatory activity. Recombinant mouse EBI3 is a non-glycosylated protein, containing 207 amino acids, with a molecular weight of 22.9 kDa.
<b>Synonyms:</b>	IL-35/EBI3, IL-27 EBI3 subunit, IL-35 EBI3 subunit
<b>Species of Origin:</b>	Mouse
<b>Expressed in:</b>	E. coli
<b>Type:</b>	Recombinant Protein
<b>Low Endotoxin:</b>	Yes

**Target Details**

<b>Gene Name:</b>	EBI3
<b>Purity/Specificity:</b>	Epstein-Barr Virus Induced Gene 3 purity was determined to be greater than 90% as determined by analysis by UV-Spectroscopy at 280nm and by reducing and non-reducing SDS-pAGE.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - O35228</a></li></ul>

## Application Details

<b>Tested Applications:</b>	SDS-PAGE
<b>Application Note:</b>	Epstein-Barr Virus Induced Gene 3 Recombinant Protein has been tested by SDS-PAGE and is suitable as a control for polyclonal or monoclonal anti-Epstein-Barr Virus Induced Gene 3 in immunological assays.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>Other:</b>	Endotoxin Level: Measured by kinetic LAL analysis and is typically $\leq 1$ EU/ $\mu$ g protein. Biologic Activity: Assay data for recombinant mouse EB13 is based upon qualitative binding to anti-EB13 antibody.


## Formulation

<b>Physical State:</b>	Lyophilized
<b>Buffer:</b>	0.1% Trifluoroacetic acid
<b>Preservative:</b>	None
<b>Stabilizer:</b>	None
<b>Reconstitution Volume:</b>	100 $\mu$ L
<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)

## Shipping & Handling

<b>Shipping Condition:</b>	Ambient
<b>Storage Condition:</b>	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
<b>Expiration:</b>	Expiration date is six (6) months from date of receipt.

## Images

**SDS-PAGE**

SDS-PAGE of Mouse Epstein-Barr Virus Induced Gene 3 Recombinant Protein. Lane 1: 1  $\mu$ g Mouse EB13 in non-reducing conditions (-). Lane 2: 1  $\mu$ g Mouse EB13 in reducing conditions (+). Lane 3: Molecular weight marker. Mouse EB13 has a predicted MW of 23.3 kDa.

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.