

**Datasheet for 010-001-S34S****KDM4B protein-GST fusion****Overview**

<b>Description:</b>	KDM4B recombinant protein-GST fusion protein - 010-001-S34S
<b>Item No.:</b>	010-001-S34S
<b>Size:</b>	20 µg
<b>Origin:</b>	Mouse
<b>Expressed in:</b>	Sf9 cells

**Product Details**

<b>Background:</b>	KDM4B or lysine (K)-specific demethylase 4B, which is also known as JMJD2B, contains a JmjN domain, a JmjC domain, a JD2H domain, 2 TUDOR domains, and a bipartite nuclear localization signal that overlaps the C-terminal part of the second TUDOR domain. KDM4B plays an essential role in human carcinogenesis through positive regulation of cyclin-dependent kinase 6 (1). KDM4B functions as a co-factor of estrogen receptor in breast cancer proliferation and mammary gland development (2). KDM4B is regulated by both ERa and HIF-1a, drives breast cancer cell proliferation in normoxia and hypoxia, and epigenetically regulates the expression of cell cycle genes such as CCND1, CCNA1, and WEE1. KDM4B Protein is ideal for investigators involved in Signaling Proteins, Deacetylase/Demethylase Proteins, Cancer, Cell Cycle, and Inflammation research.
<b>Synonyms:</b>	Jmjd2b, mKIAA0876, 4732474L06Rik, Lysine-specific demethylase 4B, EC 1.14.11.-, JmjC domain-containing histone demethylation protein 3B, Jumonji domain-containing protein 2B
<b>Species of Origin:</b>	Mouse
<b>Expressed in:</b>	Sf9 cells
<b>Type:</b>	Recombinant Protein

**Target Details**

<b>Gene Name:</b>	Kdm4b
<b>Purity/Specificity:</b>	Recombinant mouse KDM4B (1-725) was expressed by baculovirus in Sf9 insect cells using an N-Terminal Glutathione-S-Transferase fusion protein. The purity was determined to be >75% by densitometry.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">NCBI - NM_172132.2</a></li></ul>

## Application Details

<b>Application Note:</b>	KDM4B Protein is stored in 50mM Tris-HCl, pH 7.5, 50mM NaCl, 10mM glutathione, 0.1mM EDTA, 0.25mM DTT, 0.1mM PMSF, 25% glycerol. KDM4B Protein is suitable for use in Western Blot. Expect a band approximately ~125kDa on specific lysates or tissues. Specific conditions for reactivity should be optimized by the end user.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>WB:</b>	User Optimized

## Formulation

<b>Physical State:</b>	Liquid (sterile filtered)
<b>Concentration:</b>	0.1 µg/µL
<b>Buffer:</b>	See application note.

## Shipping & Handling

<b>Shipping Condition:</b>	Dry Ice
<b>Storage Condition:</b>	Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.
<b>Expiration:</b>	Expiration date is one (1) year from date of receipt.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.

