

Datasheet for 010-001-V78-0002**rMouse M-CSF Protein****Overview**

Description:	Mouse Macrophage Colony Stimulating Factor Recombinant Protein - 010-001-V78-0002
Item No.:	010-001-V78-0002
Size:	2 µg
Applications:	Cellular Assay
Origin:	Mouse
Expressed in:	E. coli

Product Details

Background:	Macrophage Colony Stimulating Factor (M-CSF) is hematopoietic growth factor produced by a wide variety of cells. M-CSF is known to stimulate differentiation of hematopoietic stem cells to monocyte-macrophage cell populations in culture. M-CSF acts through the CSF receptor 1. Although human M-CSF shows activity on mouse cells, mouse CSF shows no activity on human cells. Recombinant mouse M-CSF is a disulfide-linked homodimer, containing two 156 amino acid chains, with a total molecular weight of 36.8 kDa.
Synonyms:	MGI-IM
Species of Origin:	Mouse
Expressed in:	E. coli
Type:	Recombinant Protein
Low Endotoxin:	Yes

Target Details

Gene Name:	Csf1
Purity/Specificity:	Macrophage Colony Stimulating Factor purity was determined to be greater than 98% as determined by analysis by UV-Spectroscopy at 280nm and by reducing and non-reducing SDS-PAGE.
Relevant Links:	• UniProtKB - P07141

Application Details

Suggested Applications:	Cellular Assay (Based on references)
Application Note:	Macrophage Colony Stimulating Factor Recombinant Protein has been tested by biological activity and is suitable as a control for polyclonal or monoclonal anti-Macrophage Colony Stimulating Factor in immunological assays. Lyophilized in 10 mM sodium phosphate, 50 mM sodium chloride, pH 7.5.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	Endotoxin Level: Measured by kinetic LAL analysis and is typically ≤ 1 EU/ μ g protein. Biologic Activity: The activity is determined by dose-dependent induction of M-NFS-60 cell proliferation and is typically less than 2 ng/mL.


Formulation

Physical State:	Lyophilized
Buffer:	See application note.
Preservative:	None
Stabilizer:	None
Reconstitution Volume:	2 μ l (2-20 μ l)
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. Dilute only prior to immediate use. Maintain sterility. This product DOES NOT contain preservative. DO NOT VORTEX. We recommend adding a carrier protein such as HSA or BSA to 0.1% (i.e. 1.0 mg/mL). For best results aliquot contents and freeze at -20° C or colder. Avoid cycles of freezing and thawing. Centrifuge vial before each opening to dislodge contents from the cap and to clarify if contents are not clear after standing at room temperature.
Expiration:	Expiration date is six (6) months from date of receipt.

Images



SDS-PAGE

Bioactivity of Mouse Macrophage Colony Stimulating Factor Recombinant Protein. Serial dilutions of Mouse M-CSF, starting at 50 ng/mL, were added to NSF-60 cells. Cell proliferation was measured after 44 hours and the linear portion of the curve was used to calculate the ED50. The ED50 of Mouse M-CSF is 1.1-1.6 ng/mL. This value is comparable with the typical expected range of < 2 ng/mL.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.