

Datasheet for 100-401-A15**APC11 Antibody****Overview**

Description:	Anti-APC11 (C-terminal specific) (RABBIT) Antibody - 100-401-A15
Item No.:	100-401-A15
Size:	100 µL
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	APC11 is also known as Anaphase promoting complex subunit 11, APC11, Cyclosome subunit 11, Hepatocellular carcinoma associated RING finger protein, and HSPC214. APC11 is a component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G ₁ phase of the cell cycle. APC11 may function to recruit the E2 ubiquitin-conjugating enzymes to the complex. APC11 interacts with the cullin domain of ANAPC2 and also interacts with UBE2D2. APC11 shows both a cytoplasmic and nuclear localization. APC11 is expressed at high levels in skeletal muscle and heart; in moderate levels in brain, kidney, and liver; and at low levels in colon, thymus, spleen, small intestine, placenta, lung and peripheral blood leukocyte. APC11 is a member of the RING-type zinc finger family and is auto-ubiquitinated.
Synonyms:	rabbit anti-Apc11 Antibody, Anaphase promoting complex subunit 11 homolog antibody, APC11 anaphase promoting complex subunit 11 homolog antibody, Cyclosome subunit 11 antibody, Hepatocellular carcinoma associated RING finger protein antibody, HSPC 214 antibody, Apc
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	Antiserum

Target Details

Gene Name:	ANAPC11
Reactivity:	Human
Immunogen Type:	Conjugated Peptide

Immunogen:	This antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to amino acids 76-84 of Human APC11 (C-terminal) coupled to KLH.
Purity/Specificity:	This product is monospecific antiserum processed by delipidation and defibrination followed by sterile filtration. This product reacts with human and mouse APC11. Cross reactivity may also occur with APC11 from other sources. Sufficient sequence differences exist to suggest that this antibody would not react with other RING box proteins such as ROC1 and ROC2.
Relevant Links:	• UniProtKB - Q9NYG5 • NCBI - 19924286 • GeneID - 51529

Application Details

Application Note:	This antibody reacts with human APC11 by western blot and immunoprecipitation. The antibody immunoprecipitates in vitro translated protein and protein from overexpressing cell lysates (using HeLa and NIH-3T3, and others). Coimmunoprecipitation of related proteins (APC2) does occur. A 9.8 kDa band corresponding to human APC11 is detected. Most cell lines or tissues expressing APC11 can be used as a positive control. Researchers should determine optimal titers for other applications.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:2,000 - 1:10,000
IHC:	User Optimized
IP:	User Optimized
WB:	1:500 - 1:1,000


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	85 mg/mL by Refractometry
Buffer:	None
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Pathway

Most modifiers mature by proteolytic processing from inactive precursors (a; amino acid). Arrowheads point to the cleavage sites. Ubiquitin is expressed either as polyubiquitin or as a fusion with ribosomal proteins. Conjugation requires activating (E1) and conjugating (E2) enzymes that form thioesters (S) with the modifiers. Modification of cullins by RUB involves SCF(SKP1/cullin-1/F-box protein) /CBC(cullin-2/elongin B/elonginC) -like E3 enzymes that are also involved in ubiquitination. In contrast to ubiquitin, the UBLs do not seem to form multi-UBL chains. UCRP(ISC15) resembles two ubiquitin moieties linked head-to-tail. Whether HUB1 functions as a modifier is currently unclear. APG12 and URM1 are distinct from the other modifiers because they are unrelated in sequence to ubiquitin. Data contributed by S.Jentsch.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.