

**Datasheet for 100-4179**

## Trypsin Inhibitor Antibody

### Overview

<b>Description:</b>	Anti-Trypsin Inhibitor (RABBIT) Antibody - 100-4179
<b>Item No.:</b>	100-4179
<b>Size:</b>	2 mL
<b>Applications:</b>	WB
<b>Reactivity:</b>	Soybean
<b>Host Species:</b>	Rabbit

### Product Details

<b>Background:</b>	Soy Bean Trypsin Inhibitor is a serine protease inhibitor, an inhibitor of trypsin, therefore preventing the further breakdown of proteins.
<b>Synonyms:</b>	rabbit anti-Trypsin Inhibitor Antibody, 25 kDa trypsin inhibitor (Protease inhibitor 15 preproprotein) antibody, CRISP8 antibody, KTi antibody, Kti S antibody, KTi3 antibody, KtiS antibody, Kunitz trypsin inhibitor antibody, Kunitz type trypsin inhibitor A antibody
<b>Host Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	Antiserum

### Target Details

<b>Gene Name:</b>	KTI3
<b>Reactivity:</b>	Soybean
<b>Immunogen Type:</b>	Native Protein
<b>Immunogen:</b>	Trypsin Inhibitor [Soy Bean]
<b>Purity/Specificity:</b>	This product was prepared from monospecific antiserum by a delipidation and defibrination. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-rabbit serum, purified and partially purified Trypsin Inhibitor [Soy Bean]. Cross reactivity against Trypsin Inhibitor from other tissues and species may occur but have not been specifically determined.

**Relevant Links:**

- [UniProtKB - P01070](#)
- [NCBI - P01070.2](#)
- [GeneID - 547831](#)

## Application Details

<b>Tested Applications:</b>	WB
<b>Application Note:</b>	Anti-Trypsin Inhibitor has been tested by western blot and is suitable to be assayed against 1.0 ug of Trypsin Inhibitor [Lima Bean] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:200 to 1:1,000 of the reconstitution concentration is suggested for this product.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	1:3,000 - 1:15,000
<b>IHC:</b>	User Optimized
<b>WB:</b>	1:500 - 1:2,000

## Formulation

<b>Physical State:</b>	Lyophilized
<b>Concentration:</b>	85 mg/mL by Refractometry
<b>Buffer:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.01% (w/v) Sodium Azide
<b>Stabilizer:</b>	None
<b>Reconstitution Volume:</b>	2.0 mL
<b>Reconstitution Buffer:</b>	Restore with deionized water (or equivalent)

## Shipping & Handling

<b>Shipping Condition:</b>	Ambient
----------------------------	---------

**Storage Condition:** Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


**Expiration:** Expiration date is one (1) year from date of receipt.

## Images



### Western Blot

Rockland Rabbit anti Soybean Trypsin Inhibitor antibody was used to detect purified Soybean Trypsin Inhibitor. Samples of ~1 and 0.25 ug of purified Soybean Trypsin Inhibitor per lane were run by SDS-PAGE under reducing and non-reducing conditions and reduced samples of protein contained 4% BME and were boiled for 5 minutes. Protein was transferred to nitrocellulose and probed with Rockland Rabbit anti Soybean Trypsin Inhibitor antibody (200-4679 lot 6594 1:5K in MB-0070, ON 4 C). Primary antibody was detected with Rockland Dylight 488 conjugated Streptavidin (S000-41 Lot 21091, 1:5K) and Atto 425 conjugated goat anti rabbit (611-151-122 lot 26423C 1:10K 1.5 hr RT in MB-070) and imaged on the BioRad VersaDoc imaging system.



### Western Blot

Rockland Biotin conjugated Rabbit anti-trypsin inhibitor antibody (200-4679 lot 6594) and Rabbit anti streptavidin (200-4195 lot 23495) were used to detect target proteins Trypsin Inhibitor (left) and Streptavidin (right) under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified target proteins contained 4% BME and were boiled for 5 minutes. Samples of ~1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:1000 dilution of primary antibody (ON 4 C). Detection shown was using Dylight 649 conjugated Donkey anti rabbit (611-743-127 lot 20831 1:10K 1.5 hr RT in MB-070) and imaged on the BioRad VersaDoc System

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.