

**Datasheet for 200-101-037S**

## Glucose Oxidase Antibody

### Overview

<b>Description:</b>	Anti-Glucose Oxidase (GOAT) Antibody - 200-101-037S
<b>Item No.:</b>	200-101-037S
<b>Size:</b>	25 µL
<b>Applications:</b>	WB
<b>Reactivity:</b>	Aspergillus niger
<b>Host Species:</b>	Goat

### Product Details

<b>Background:</b>	Glucose Oxidase enzyme is widely applied for the determination of glucose in body fluids and in removing residual glucose or oxygen from foods and beverages. Furthermore, glucose oxidase-producing molds such as aspergillus and penicillium species are used for the biological production of gluconic acid.
<b>Synonyms:</b>	goat anti-Glucose Oxidase Antibody, Beta D Glucose Oxygen 1 Oxido Reductase antibody, Glucose oxidase (Precursor) antibody, Glucose Oxyhydrase antibody, GOD antibody
<b>Host Species:</b>	Goat
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	IgG

### Target Details

<b>Gene Name:</b>	gox
<b>Reactivity:</b>	Aspergillus niger
<b>Immunogen Type:</b>	Native Protein
<b>Immunogen:</b>	Glucose Oxidase [Aspergillus niger]

**Purity/Specificity:** Anti-Glucose Oxidase Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Glucose Oxidase [*Aspergillus niger*]. Cross reactivity against Glucose Oxidase from other tissues and species may occur but have not been specifically determined.

**Relevant Links:**

- [UniProtKB - P13006](#)
- [NCBI - P13006.1](#)

## Application Details

<b>Tested Applications:</b>	WB
<b>Application Note:</b>	Anti-Glucose Oxidase has been tested in western blot and is suitable for use in ELISA and immunohistochemistry. Specific conditions for reactivity should be optimized by the end user.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	1:2,000 - 1:10,000
<b>IHC:</b>	User Optimized
<b>WB:</b>	1:1000 - 1:5000

## Formulation

<b>Physical State:</b>	Liquid (sterile filtered)
<b>Concentration:</b>	1.0 mg/ml by UV absorbance at 280 nm
<b>Buffer:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.01% (w/v) Sodium Azide
<b>Stabilizer:</b>	None



## Shipping & Handling

<b>Shipping Condition:</b>	Dry Ice
----------------------------	---------

**Storage Condition:** Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.

**Expiration:** Expiration date is three (3) months from date of receipt.

## Images



### Western Blot

Rockland Goat anti Glucose Oxidase antibody was used to detect Glucose Oxidase under reducing and non-reducing conditions. Reduced samples of purified Glucose Oxidase contained 4% BME and were boiled for 5 minutes. Samples of ~1 and 0.25 ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with Rockland Goat anti-Glucose Oxidase antibody (p/n 200-1137, lot 165) at 1:5K in MB-070, ON 4°C. Primary antibody was detected with Rockland Dylight 649 conjugated Donkey anti-Goat (p/n 605-743-125, lot 20834) at 1:10K 1.5 hr RT in MB-070 and imaged on the BioRad VersaDoc imaging system.

### Western Blot

Western Blot of Goat anti-Glucose Oxidase antibody. Lane 1: Glucose Oxidase. Load: 50 ng per lane. Primary antibody: Glucose Oxidase antibody at 1:1,000 for overnight at 4°C. Secondary antibody: Peroxidase goat secondary antibody (p/n 605-4302) at 1:40,000 for 30 min at RT. Block: (p/n MB-070) for 30 min at RT. Predicted/Observed size: 65-70 kDa, 65-70 kDa for Glucose Oxidase. Other band(s): None.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.