

Datasheet for 200-101-093S**Sarcosine Oxidase Antibody****Overview**

Description:	Anti-Sarcosine Oxidase (Microbial) (GOAT) Antibody - 200-101-093S
Item No.:	200-101-093S
Size:	25 µL
Applications:	Dot Blot, WB
Reactivity:	Sarcosine Oxidase (Microbial)
Host Species:	Goat

Product Details

Background:	Sarcosine Oxidase catalyzes the oxidative demethylation of sarcosine to yield glycine, hydrogen peroxide and 5,10-methylenetetrahydrofolate.
Synonyms:	goat anti-Sarcosine Oxidase Antibody, Sarcosine oxidase subunit alpha, Sarcosine oxidase subunit
Host Species:	Goat
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	soxA
Reactivity:	Sarcosine Oxidase (Microbial)
Immunogen Type:	Native Protein
Immunogen:	Sarcosine Oxidase [Microbial]
Purity/Specificity:	Anti-SARCOSINE OXIDASE is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Sarcosine Oxidase [Microbial]. Cross reactivity against Sarcosine Oxidase from other sources is unknown.

Relevant Links:

- [UniProtKB - Q46337](#)
- [NCBI - YP_004446819.1](#)
- [GeneID - 10586229](#)

Application Details

Tested Applications:	Dot Blot, WB
Application Note:	Anti-Sarcosine Oxidase Antibody has been tested by dot blot and western blot and is suitable for ELISA assay against 1.0 ug of Sarcosine Oxidase [Microbial] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) code #605-4302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:12,000 of the reconstitution concentration is suggested for this product.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000 - 1:20,000
IP:	1:100
WB:	1:500 - 1:5,000

Formulation


Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/ml by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Rockland Goat anti Sarcosine Oxidase antibody was used to detect Sarcosine Oxidase. Samples of ~1 and 0.25 ug of purified Sarcosine Oxidase per lane were run by SDS-PAGE under reducing and non-reducing conditions and reduced samples contained 4% BME and were boiled for 5 minutes. Protein was transferred to nitrocellulose and probed with Rockland Goat anti Sarcosine Oxidase (200-101-093 lot 5951 1:5K in MB-0070, ON 4 C). Primary antibody was detected with Rockland Dylight 649 conjugated Donkey anti Goat (605-743-125 1:10K 1.5 hr RT in MB-070 and imaged on the BioRad VersaDoc imaging system.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.