

Datasheet for 200-101-231-0100**Sucrose Phosphorylase Antibody****Overview**

Description:	Anti-Sucrose Phosphorylase (E. coli) (GOAT) Antibody - 200-101-231-0100
Item No.:	200-101-231-0100
Size:	100 µg
Applications:	WB
Reactivity:	E. coli
Host Species:	Goat

Product Details

Background:	Sucrose phosphorylase catalyzes as Sucrose + phosphate = D-fructose + alpha-D-glucose 1-phosphate.
Synonyms:	goat anti-Sucrose Phosphorylase Antibody, Putative sucrose phosphorylase, Sucrose glucosyltransferase
Host Species:	Goat
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	ycjM
Reactivity:	E. coli
Immunogen Type:	Native Protein
Immunogen:	Sucrose Phosphorylase [E.coli]
Purity/Specificity:	Anti-Sucrose phosphorylase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Sucrose Phosphorylase [E.coli]. Cross reactivity against Sucrose Phosphorylase from other sources is unknown.

Relevant Links:	• UniProtKB - P76041 • NCBI - YP_852475.1 • GeneID - 4492194
------------------------	--

Application Details

Tested Applications:	WB
Application Note:	Anti-Sucrose phosphorylase Antibody has been tested by western blot and is suitable to be assayed against 1.0 µg of Sucrose Phosphorylase [E.coli] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:6,500 to 1:32,000 of the reconstitution concentration is suggested for this product.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000 - 1:20,000
IP:	1:100
WB:	1:500 - 1:5,000

Formulation

Physical State:	Lyophilized
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None
Reconstitution Volume:	100 µL
Reconstitution Buffer:	Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition:	Ambient
----------------------------	---------

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Rockland Goat anti Sucrose Phosphorylase antibody (200-101-231 lot 7851) was used to detect purified Sucrose Phosphorylase under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified protein contained 4% BME and were boiled for 5 minutes. Samples of ~1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:3000 dilution of primary antibody (ON 4 C in MB-070). Detection shown was using Dylight 488 conjugated Donkey anti goat (605-741-125 lot 21094 1:10K in TBS/MB-070 1 hr RT). Images were collected using the BioRad VersaDoc System



Western Blot

Western Blot of Goat Anti-Sucrose Phosphorylase Antibody. Lane 1: Opal Prestained Ladder (p/n MB-210-0500). Lane 2: HeLa Lysate (10µg) spiked with Sucrose Phosphorylase protein (50ng). Lane 3: HeLa Lysate (10µg) spiked with Sucrose Phosphorylase protein (20ng). Lane 4: HeLa Lysate (10µg) spiked with Sucrose Phosphorylase protein (10ng). Lane 5: HeLa Lysate (10µg) (p/n W09-000-364). Primary Antibody: Anti-Sucrose Phosphorylase at 1µg/mL overnight at 2-8°C. Secondary Antibody: Donkey Anti-Goat IgG HRP mx8 custom p/n at 1:40,000 for 30mins at RT. Block: BlockOut Buffer (p/n MB-073). Expect/Observe MW: ~50kDa.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.