

Datasheet for 200-103-212**Bilirubin Oxidase Antibody Peroxidase Conjugated****Overview**

Description:	Anti-Bilirubin Oxidase (Myrothecium verrucaria) (GOAT) Antibody Peroxidase Conjugated (BULK ORDER) - 200-103-212
Item No.:	200-103-212
Size:	20 mg
Applications:	WB
Reactivity:	Myrothecium verrucaria
Host Species:	Goat

Product Details

Background:	Bilirubin Oxidase is key in the oxidation of bilirubin and other tetrapyrroles. This enzyme belongs to the family of oxidoreductases, to be specific those acting on the CH-CH group of donor with oxygen as acceptor. The systematic name of this enzyme class is bilirubin:oxygen oxidoreductase. This enzyme is also called bilirubin oxidase M-1. This enzyme participates in porphyrin and chlorophyll metabolism.
Synonyms:	goat anti-Bilirubin Oxidase HRP conjugation, goat anti-Bilirubin Oxidase Antibody Peroxidase Conjugated
Host Species:	Goat
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	BLRO_MYRVE
Reactivity:	Myrothecium verrucaria
Immunogen Type:	Native Protein
Immunogen:	Bilirubin Oxidase [Myrothecium verrucaria]

Purity/Specificity:	Bilirubin Oxidase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum as well as purified and partially purified Bilirubin Oxidase [Myrothecium verrucaria]. Cross reactivity against Bilirubin Oxidase from other sources is unknown.
----------------------------	---

Relevant Links:	• UniProtKB - Q12737
------------------------	--

Application Details

Tested Applications:	WB
Application Note:	Anti-Bilirubin Oxidase has been tested by western blot and is suitable to be assayed against 1.0 ug of Bilirubin Oxidase [Myrothecium verrucaria] in a standard capture ELISA using ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:500 to 1:2,500 of the reconstitution concentration is suggested for this product.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000 - 1:20,000
IP:	1:100
WB:	1:500 - 1:5,000

Formulation

Physical State:	Lyophilized
Concentration:	10 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	2.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)



Shipping & Handling

Shipping Condition:	Ambient
----------------------------	---------

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Rockland Goat anti Bilirubin Oxidase antibody (200-101-212 lot 6533) was used to detect Bilirubin Oxidase under reducing (R) conditions. A samples of ~1ug of purified protein contained 4% BME and was boiled for 5 minutes and run by SDS-PAGE on a 4-20% acrylamide gel. Protein was transferred to nitrocellulose and probed with 1:3000 dilution of primary antibody (ON 4 C in MB-070). Detection shown was using Dylight 488 conjugated Donkey anti goat (605-741-125 lot 21094 1:10K in TBS/MB-070 1 hr RT) and Dylight 649 conjugated Donkey anti goat (605-743-125 lot 20834 1:10K in TBS/MB-070) 1 hr RT. Images were collected using the BioRad VersaDoc System

Western Blot

Western Blot of Goat anti-Bilirubin Oxidase (Myrothecium verrucaria) Antibody Peroxidase Conjugated. Lane 1: Bilirubin Oxidase (Myrothecium verrucaria). Load: 50 ng per lane. Primary antibody: Goat anti-Bilirubin Oxidase (Myrothecium verrucaria) Antibody Peroxidase Conjugated at 1:1,000 overnight at 4°C. Secondary antibody: n/a. Block: MB-070 for 30 minutes at RT. Predicted/Observed size: 64 kDa, 60 kDa for Bilirubin Oxidase. Other band(s): Bilirubin Oxidase splice variants and isoforms.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.