Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 200-106-237

Beta-Phosphoglucomutase Antibody Biotin Conjugated

Overview

Description:	Anti-Beta Phosphoglucomutase (GOAT) Antibody Biotin Conjugated (BULK ORDER) - 200-106-237
Item No.:	200-106-237
Size:	10 mg
Applications:	ELISA, WB
Reactivity:	Lactococcus lacti
Host Species:	Goat

Product Details

Background:	Anti-Beta-Phosphoglucomutase antibody is located in the cytoplasm, working to catalyze the transference of a phosphoryl group on a glucose or lactose monomer from the 1' to the 6' position in the forward direction or 6' to the 1' position in the reverse direction. Anti-Beta-Phosphoglucomutase is essential to regulating the flow of carbohydrate intermediates in glycolysis as well as the formation of the sugar nucleotide UDP-glucose. Part of the HAD-like hydrolase superfamily, Anti-Beta-Phosphoglucomutase antibody is ideal for researchers interested in Metabolism research.
Synonyms:	goat anti-Beta-Phosphoglucomutase Antibody biotin Conjugation, biotin conjugated goat anti-Beta PGM antibody, Glucose phosphomutase 1 antibody, Glucose phosphomutase antibody, PGM antibody, pgmB antibody
Host Species:	Goat
Conjugate:	Biotin
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	pgmB
Reactivity:	Lactococcus lacti
Immunogen Type:	Native Protein

Immunogen:	beta Phosphoglucomutase [Lactococcus lacti]
Purity/Specificity:	Anti-Beta-Phosphoglucomutase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Goat Serum as well as purified and partially purified Beta-Phosphoglucomutase [Lactococcus lacti]. Cross reactivity against Beta-Phosphoglucomutase from other sources is unknown.
Relevant Links:	• UniProtKB - P71447 • NCBI - CAA94734.1 • GeneID - 1114041

Application Details

Tested Applications:	ELISA, WB
Application Note:	Anti-Beta-Phosphoglucomutase Biotin has been tested by ELISA and western blot. This product is assayed against 1.0 ug of Beta-Phosphoglucomutase in a standard capture ELISA using Peroxidase Conjugated Streptavidin #S000-03 and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:50,000 is suggested for this product.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000 - 1:25,000
WB:	1:500 - 1:3,000


Formulation

Physical State:	Lyophilized
Concentration:	10.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot
Rockland Goat anti antibody (200-101-237 lot 8180) was used to detect purified Beta Phospho Glucomutase under reducing (R) and non-reducing (NR) conditions. Reduced samples of protein contained 4% BME and were boiled for 5 minutes. Samples of ~1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:3000 dilution of primary antibody (ON 4 C in MB-070). Detection shown was using Dylight 488 conjugated Donkey anti goat (605-741-125 lot 21094 1:10K in TBS/MB-070 1 hr RT). Images were collected using the BioRad VersaDoc System.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.