

Datasheet for 200-1376**Pepsin Antibody Peroxidase Conjugated****Overview**

Description:	Anti-Pepsin (GOAT) Antibody Peroxidase Conjugated (BULK ORDER) - 200-1376
Item No.:	200-1376
Size:	20 mg
Applications:	Dot Blot, ELISA, WB
Reactivity:	Pig
Host Species:	Goat

Product Details

Background:	Pepsin is one of three principal protein-degrading, or proteolytic, enzymes in the digestive system, the other two being chymotrypsin and trypsin. Pepsin is commonly used in the preparation produce either an Fab or an F(ab') ₂ fragment from antibodies.
Synonyms:	goat anti-Pepsin Antibody HRP Conjugated, Peroxidase Conjugated goat anti-Pepsin Antibody, Pepsin A antibody, Pepsin a precursor antibody, Pepsin antibody, Pepsinogen 1 second locus antibody, Pepsinogen 3 group 1 antibody, Pepsinogen a3 antibody
Host Species:	Goat
Conjugate:	Peroxidase (HRP)
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	PGA
Reactivity:	Pig
Immunogen Type:	Native Protein
Immunogen:	Pepsin [Porcine Stomach]

Purity/Specificity: Pepsin is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Goat Serum as well as purified and partially purified Pepsin [Porcine Stomach]. Cross reactivity against Pepsin from other tissues and species may occur but have not been specifically determined.

Relevant Links:

- [UniProtKB - P00791](#)
- [NCBI - AAA31095.1](#)
- [GeneID - 396892](#)

Application Details

Tested Applications:	Dot Blot, ELISA, WB
Application Note:	Anti-Pepsin peroxidase has been tested by ELISA, dot blot, and western blot and is useful in immunohistochemistry. Optimal titers should be obtained by researchers if not provided.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:75,000
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:10,000

Formulation

Physical State:	Lyophilized
Concentration:	10.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	2.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition:	Ambient
----------------------------	---------

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Peroxidase conjugated Goat anti-Pepsin antibody. Lane 1: Pepsin. Lane 2: none. Load: 100 ng per lane. Primary antibody: none. Secondary antibody: Peroxidase goat pepsin antibody at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 35 kDa for Pepsin. Other band(s): none.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.