

**Datasheet for 200-301-200****GST Antibody****Overview**

<b>Description:</b>	Anti-GST (MOUSE) Monoclonal Antibody - 200-301-200
<b>Item No.:</b>	200-301-200
<b>Size:</b>	500 µg
<b>Applications:</b>	ELISA, WB, IP
<b>Reactivity:</b>	GST-Tag
<b>Host Species:</b>	Mouse

**Product Details**

<b>Background:</b>	Rockland produces a wide range of GST antibodies in our laboratories. Select GST antibodies from several monoclonal and/or polyclonal GST antibodies listed below. Select appropriate GST antibodies for your research by isotype, epitope, applications and species reactivity. GST (Glutathione-S-Transferase) is a protein expression tag commonly used in molecular biology. Anti-GST will react with synthetic construct present in most known GST containing cloning or expression vectors. GST is responsible for the conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. The amino acid sequence GST is highly conserved in most organisms including mammals. GST exists as a 26 kDa homodimer.
<b>Synonyms:</b>	mouse anti-GST antibody, Glutathione-S-Transferase, Anti-GST monoclonal antibody
<b>Host Species:</b>	Mouse
<b>Clonality:</b>	Monoclonal
<b>Clone ID:</b>	3D4
<b>Format:</b>	IgG1

**Target Details**

<b>Reactivity:</b>	GST-Tag
<b>Immunogen Type:</b>	Native Protein
<b>Immunogen:</b>	A BALB/c mouse was immunized with Glutathione-S-Transferase [Schistosoma japonicum].

**Purity/Specificity:** Anti-GST monoclonal antibody was purified from concentrated tissue culture supernate by Protein A chromatography which reacts specifically with GST. Cross reactivity against Glutathione-S-Transferase from other sources may occur but has not been specifically determined.

---

## Application Details

**Tested Applications:** ELISA, WB

**Suggested Applications:** IP (Based on references)

**Application Note:** Anti-GST antibody has been tested by ELISA and western blot and is suitable for most immunological techniques requiring high titer binding and lot-to-lot consistency.

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

**ELISA:** 1:50,000 - 1:150,000

**IHC:** User Optimized

**IP:** User Optimized

**WB:** 1:1,000 - 1:10,000

---

## Formulation

**Physical State:** Liquid (sterile filtered)

**Concentration:** 1.006 mg/mL by UV absorbance at 280 nm

**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Preservative:** 0.01% (w/v) Sodium Azide

**Stabilizer:** None

---

## Shipping & Handling

**Shipping Condition:** Dry Ice

**Storage Condition:** Store anti-GST at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Expiration:** Expiration date is one (1) year from date of receipt.

---

## Images



### Western Blot

Western Blot of anti-GST antibody. Lane 1: GST recombinant protein (p/n 000-001-200). Lane 2: HeLa cell lysates expressing recombinant GST protein (p/n W09-000-364/000-001-200). Load: 0.1 µg per lane. Primary antibody: GST antibody at 1.0µg/ml for 1 h at room temperature. Secondary antibody: IRDye800™ goat anti-mouse secondary antibody at 1:5,000 for 45 min at RT. Block: 5% BLOTTO (p/n B501-0500) overnight at 4°C. Other band(s): none.



### Western Blot

Western Blot of Mouse-anti-GST antibody. Lane 1: molecular weight marker. Lane 2: Mouse-anti-GST monoclonal antibody (blue), Rabbit anti-Transferrin (green), and Goat-anti-Alpha-1-Anti-Trypsin (red) were used in a multiplex system to detect target proteins under reducing conditions in albumin depleted human serum with 320 ng of added GST. Load: 1 µg per lane. Primary antibody: Each primary antibody at 1:1000 for overnight at 4°C. Secondary antibody: Donkey anti-Rabbit IgG DyLight549 (p/n 611-742-127), Donkey anti-Mouse IgG DyLight 488 (p/n 610-741-002), and Donkey anti-Goat IgG DyLight 649 (p/n 605-743-002) secondary antibody at 1:10,000 for 30 min at RT. Block: MB-070 overnight at 4°C.

## References

- Foss S et al. Human IgG Fc-engineering for enhanced plasma half-life, mucosal distribution and killing of cancer cells and bacteria. *Nat Commun.* (2024)
- Yoshida, Y et al. Elevation of autoantibody level against PDCD11 in patients with transient ischemic attack. *Oncotarget* (2018)
- Bolch, SN et al. A Splice Variant of Bardet-Biedl Syndrome 5 (BBS5) Protein that Is Selectively Expressed in Retina. *PLoS One* (2016)
- Smith, TS et al. Light-dependent phosphorylation of Bardet-Biedl syndrome 5 in photoreceptor cells modulates its interaction with arrestin1. *Cellular and Molecular Life Sciences : Cmls* (2013)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.