

Datasheet for 200-301-279**LEFTY A Antibody****Overview**

Description:	Anti-LEFTY A (MOUSE) Monoclonal Antibody - 200-301-279
Item No.:	200-301-279
Size:	100 µg
Applications:	ELISA, WB
Reactivity:	Human
Host Species:	Mouse

Product Details

Background:	During vertebrate embryogenesis, a left-right axis is established. Secreted growth factors of the TGF-beta family, including gene products derived from nodal, lefty-1 and lefty-2, play crucial roles in establishing left-right asymmetries. TGF-beta (Transforming growth factor-beta) is a pleiotropic cytokine that regulates growth and differentiation of diverse types of cells. TGF-beta actions are directed by ligand-induced activation of TGF-beta receptors. Complexes formed move into the nucleus, where they act as components of a transcriptional complex. Lefty, a novel member of the TGF-beta superfamily, inhibits TGF-beta signaling. Lefty acts to inhibit phosphorylation of Smad2 following activation of the TGF-beta receptor. Lefty also inhibits events downstream from R-Smad phosphorylation. Lefty provides a repressed state of TGF-beta-responsive genes. The Lefty family is comprised of Lefty 1 and Lefty 2 in mouse, and Lefty A and Lefty B in humans. Members of the TGF-beta superfamily require processing for their activation. Cleavage is therefore an essential step for Lefty activation. Lefty is synthesized as a large inactive precursor (42 kDa) that must be endoproteolytically processed to release the bioactive polypeptide (28 kDa and 34 kDa forms). The 28kDa form induces MAPK activity.
Synonyms:	mouse anti-leftyA antibody, mouse anti-protein lefty 2 antibody, Left-right determination factor 2 antibody, Protein lefty-2 antibody, Left-right determination factor A antibody, Protein lefty-A antibody, Transforming growth factor beta-4 antibody, TGF-beta-4 antibody, Endometrial bleeding-associated factor antibody
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	7C5G1H6H10
Format:	IgG2a

Target Details

Gene Name:	LEFTY2
Reactivity:	Human
Immunogen Type:	Recombinant Protein
Immunogen:	A BALB/c mouse was immunized with a recombinant form of 6X HIS tagged human LEFTY.
Purity/Specificity:	This protein A purified mouse monoclonal antibody reacts with a 30 kDa protein corresponding to human LEFTY. No reactivity occurs against 6X HIS tag. Significant sequence homology exists between the human and mouse forms of LEFTY. Cross reactivity is expected with mouse and rat forms of the protein. Reactivity with other sources has not been determined.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - O00292• NCBI - NP_001165896.1• GenelD - 7044

Application Details

Tested Applications:	ELISA, WB
Application Note:	This antibody has been tested by ELISA and western blotting. The antibody may be used for other applications, such as RIA, immunohistochemistry or immunoprecipitation, but specific reaction conditions have not been developed.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000
WB:	1:2,000


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.051 mg/mL by nanodrop at 205 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Mab anti-Human LEFTY antibody (clone 7C5G1H6H10) is shown to detect by western blot partially purified recombinant 6X His tagged human LEFTY. Lane 1 contains an unrelated 6X His tagged protein and shows that the antibody does not recognize the epitope tag. Lane 2 contains partially purified recombinant human LEFTY. Detection occurs after 1.0 µg of protein is loaded in each lane. The blot was incubated with a 1:2,000 dilution of Mab anti-Human LEFTY at room temperature for 30 min followed by detection using IRDye™800 labeled Goat-a-Mouse IgG [H&L] (610-132-121) diluted 1:1,000. The antibody may be used to detect endogenous human LEFTY. IRDye™800 fluorescence image was captured using the Odyssey® Infrared Imaging System developed by LI-COR. IRDye is a trademark of LI-COR, Inc. Other detection systems will yield similar results.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.