

Datasheet for 200-301-880**Alpha-Tubulin Antibody****Overview**

Description:	Anti-Alpha-Tubulin (MOUSE) Monoclonal Antibody - 200-301-880
Item No.:	200-301-880
Size:	100 µg
Applications:	ELISA, IF, IHC, Multiplex, WB
Reactivity:	Human
Host Species:	Mouse

Product Details

Background:	Microtubules are involved in a wide variety of cellular activities ranging from mitosis and transport events to cell movement and the maintenance of cell shape. Tubulin itself is a globular protein consisting of two polypeptides (alpha and beta tubulin). Alpha and beta tubulin dimers are assembled to 13 protofilaments that form a microtubule of 22-nm diameter. Tyrosine ligase adds a C-terminal tyrosine to monomeric alpha tubulin. Assembled microtubules can again be detyrosinated by a cytoskeleton-associated carboxypeptidase. Detyrosinated alpha tubulin is referred to as Glu-tubulin. Another post-translational modification of detyrosinated alpha tubulin is C-terminal polyglutamylation, which is characteristic of microtubules in neuronal cells and the mitotic spindle. This antibody makes an excellent loading control.
Synonyms:	mouse anti-Alpha-Tubulin Antibody, Tubulin alpha-1B chain, Tubulin alpha-ubiquitous chain, Alpha-tubulin ubiquitous, Tubulin K-alpha-1, TUBA1B, tubulin loading control
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	17H11.F10
Format:	IgG1

Target Details

Gene Name:	TUBA1B
Reactivity:	Human

Immunogen:	Anti-Tubulin Loading Control Antibody was produced by repeated immunizations with a synthetic peptide corresponding to residues near the C terminal end of human alpha tubulin protein.
Purity/Specificity:	Anti-Tubulin Loading Control Antibody was purified by Protein A chromatography. This Loading Control antibody is directed against alpha tubulin. A BLAST analysis was used to suggest antibody reactivity with alpha tubulin from a wide range of organisms, including avian, mammalian aquatic, parasitic and alga sources based on 100% homology for the immunogen sequence. Cross reactivity will occur with all isoforms of alpha tubulin. Such broad reactivity makes this antibody useful as an excellent loading control.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P68363• NCBI - 17986283

Application Details

Tested Applications:	ELISA, IF, IHC, Multiplex, WB
Application Note:	Anti-Tubulin Antibody has been tested for use in ELISA, immunohistochemistry, immunofluorescence microscopy and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band at ~50 kDa in size corresponding to alpha tubulin by western blotting in most cell lysates or extracts.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:300,000
IF:	0.1 µg/mL
IHC:	2.5 µg/mL
WB:	1:1,000


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store Anti-Tubulin Antibody at -20° C prior to opening. Aliquot Loading Control Antibody and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge Tubulin Loading Control Antibody if not completely clear after standing at room temperature. This Anti-Tubulin Loading Control Antibody is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images




Western Blot

miR-155 and FOXP3 down regulate endogenous ZEB2 in human breast cancer cells resulting in altered levels of EMT markers Vimentin and E-cadherin(A) Relative abundance of ZEB2 and ZEB1 protein in WT, GFP or FOXP3 overexpressing BT549 cells transfected with miR-155 or miR-control. Relative abundance of protein was determined by quantitation of the abundance of ZEB2 or ZEB1 proteins normalised to reference protein α -Tubulin by western blot analysis. Quantitation of bands was carried out using Image J software. Mean + SD plotted. Student's t test ***P < 0.001. ZEB1 protein expression as above. n = 3 experiments. (B) ZEB2 and ZEB1 protein in WT, GFP or FOXP3 overexpressing BT549 cells transfected with miR-155 or miR-control by western blot. Representative western blot shown. (C) Relative abundance of Vimentin and E-cadherin protein in WT, GFP or FOXP3 overexpressing BT549 cells transfected with miR-155 or miR-control. Relative abundance of protein was determined by quantitating the abundance of E-cadherin or Vimentin proteins and normalising to reference protein β -Actin by western blot analysis. Quantitation of bands was carried out using Image J software. Mean + SD plotted. Student's t test ***P < 0.001, **P < 0.01. n = 3 experiments. (D) Vimentin and E-cadherin protein in WT, GFP or FOXP3 overexpressing BT549 cells transfected with miR-155 or miR-control analysed by western blot. Representative western blot shown. Figure provided by CiteAb. Source: Oncotarget, PMID: 29963231.

Western Blot


Smad3 is required for TGF- β to regulate expression of Fmod and Adamtsl2. (A) Primary sclerotome was treated with vehicle control or TGF β 1 for 2 h, 8 h or 24 h. Immunoblot was used to determine the levels of pSmad3, pSmad2 and




total Smad2/3. α -Tubulin was used as a general loading control. (B) Quantification of pSmad3 and (C) pSmad2 levels relative to total Smad2/3 are shown. Immunoblots were quantified using ImageJ. (*) indicates significance, $p < 0.05$, $n = 3$ for each (D) Sclerotome was pretreated with vehicle control or SIS3 (5 μ M and 10 μ M) for 24 h and then treated with or without TGF β 1 for 8 h. pSmad3 and Smad2/3 protein levels were analyzed via immunoblot. α -Tubulin was used as a general loading control. (E) Smad3 activity was quantified from the immunoblots using ImageJ as the relative levels of pSmad3 normalized to total Smad2/3. (*) indicates significance, $p < 0.05$, $n = 3$ (F) Relative levels of Scx, Fmod, Adamts12 and Prg4 mRNA were determined by qPCR after indicated treatment with SIS3. All mRNA levels were normalized to the housekeeping gene Hprt and then analyzed for significance using REST (*). Results are shown relative to TGF β 1 treated cells. $n = 3$ (G) Cells were infected with AdDNSmad2 or AdGFP control virus 48 h before cells were treated with or without TGF β 1 for 8 h. Immunoblot was used to visualize relative levels of pSmad3, pSmad2 and total Smad2/3. α -Tubulin is shown as a general loading control. (H) pSmad3 and (I) pSmad2 activity were quantified using ImageJ as the level of each phosphoSmad normalized to total Smad2/3. (*) indicates significance, $p < 0.05$, $n = 3$ (J) RNA was extracted from cells that were infected with AdDNSmad2 or AdGFP and treated or untreated with TGF β 1. Relative levels of Scx and Adamts12 mRNA were determined by qPCR. Expression was normalized to the housekeeping gene Hprt. Results are shown relative to the untreated control (*) indicates significance, $p < 0.05$, $n = 5$ (K) Cells were infected with AdSmad3 or AdGFP for 48 h and Smad3 expression was verified via immunoblot. (L) Immunoblots were quantified using ImageJ. (*) indicates significance, $p < 0.05$, $n = 3$ (M) RNA was isolated from cells infected with AdSmad3 or AdGFP for 48 h and then treated or untreated with TGF β 1 for 8 h. Relative levels of Scx, Fmod, Adamts12, and Prg4 mRNA were determined by qPCR. Results are shown relative to Ad-GFP infected, untreated controls. All mRNA levels were normalized to the housekeeping gene Hprt. (*) indicates significance, $p < 0.05$, $n = 3$. Detailed results from qPCR REST analysis are shown in Tables S4–S6. Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.

Western Blot

Smad3 is required for TGF- β to regulate expression of Fmod




and Adamtsl2. (A) Primary sclerotome was treated with vehicle control or TGF β 1 for 2 h, 8 h or 24 h. Immunoblot was used to determine the levels of pSmad3, pSmad2 and total Smad2/3. α -Tubulin was used as a general loading control. (B) Quantification of pSmad3 and (C) pSmad2 levels relative to total Smad2/3 are shown. Immunoblots were quantified using ImageJ. (*) indicates significance, $p < 0.05$, $n = 3$ for each (D) Sclerotome was pretreated with vehicle control or SIS3 (5 μ M and 10 μ M) for 24 h and then treated with or without TGF β 1 for 8 h. pSmad3 and Smad2/3 protein levels were analyzed via immunoblot. α -Tubulin was used as a general loading control. (E) Smad3 activity was quantified from the immunoblots using ImageJ as the relative levels of pSmad3 normalized to total Smad2/3. (*) indicates significance, $p < 0.05$, $n = 3$ (F) Relative levels of Scx, Fmod, Adamts2 and Prg4 mRNA were determined by qPCR after indicated treatment with SIS3. All mRNA levels were normalized to the housekeeping gene Hprt and then analyzed for significance using REST (*). Results are shown relative to TGF β 1 treated cells. $n = 3$ (G) Cells were infected with AdDNSmad2 or AdGFP control virus 48 h before cells were treated with or without TGF β 1 for 8 h. Immunoblot was used to visualize relative levels of pSmad3, pSmad2 and total Smad2/3. α -Tubulin is shown as a general loading control. (H) pSmad3 and (I) pSmad2 activity were quantified using ImageJ as the level of each phosphoSmad normalized to total Smad2/3. (*) indicates significance, $p < 0.05$, $n = 3$ (J) RNA was extracted from cells that were infected with AdDNSmad2 or AdGFP and treated or untreated with TGF β 1. Relative levels of Scx and Adamtsl2 mRNA were determined by qPCR. Expression was normalized to the housekeeping gene Hprt. Results are shown relative to the untreated control (*) indicates significance, $p < 0.05$, $n = 5$ (K) Cells were infected with AdSmad3 or AdGFP for 48 h and Smad3 expression was verified via immunoblot. (L) Immunoblots were quantified using ImageJ. (*) indicates significance, $p < 0.05$, $n = 3$ (M) RNA was isolated from cells infected with AdSmad3 or AdGFP for 48 h and then treated or untreated with TGF β 1 for 8 h. Relative levels of Scx, Fmod, Adamtsl2, and Prg4 mRNA were determined by qPCR. Results are shown relative to Ad-GFP infected, untreated controls. All mRNA levels were normalized to the housekeeping gene Hprt. (*) indicates significance, $p < 0.05$, $n = 3$. Detailed results from qPCR REST analysis are shown in Tables S4–S6. Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.



Western Blot

Smad3 is required for TGF-β to regulate expression of Fmod and Adamtsl2. (A) Primary sclerotome was treated with vehicle control or TGFβ1 for 2 h, 8 h or 24 h. Immunoblot was used to determine the levels of pSmad3, pSmad2 and total Smad2/3. α-Tubulin was used as a general loading control. (B) Quantification of pSmad3 and (C) pSmad2 levels relative to total Smad2/3 are shown. Immunoblots were quantified using ImageJ. (*) indicates significance, $p < 0.05$, $n = 3$ for each (D) Sclerotome was pretreated with vehicle control or SIS3 (5 μM and 10 μM) for 24 h and then treated with or without TGFβ1 for 8 h. pSmad3 and Smad2/3 protein levels were analyzed via immunoblot. α-Tubulin was used as a general loading control. (E) Smad3 activity was quantified from the immunoblots using ImageJ as the relative levels of pSmad3 normalized to total Smad2/3. (*) indicates significance, $p < 0.05$, $n = 3$ (F) Relative levels of Scx, Fmod, Adamtsl2 and Prg4 mRNA were determined by qPCR after indicated treatment with SIS3. All mRNA levels were normalized to the housekeeping gene Hprt and then analyzed for significance using REST (*). Results are shown relative to TGFβ1 treated cells. $n = 3$ (G) Cells were infected with AdDNSmad2 or AdGFP control virus 48 h before cells were treated with or without TGFβ1 for 8 h. Immunoblot was used to visualize relative levels of pSmad3, pSmad2 and total Smad2/3. α-Tubulin is shown as a general loading control. (H) pSmad3 and (I) pSmad2 activity were quantified using ImageJ as the level of each phosphoSmad normalized to total Smad2/3. (*) indicates significance, $p < 0.05$, $n = 3$ (J) RNA was extracted from cells that were infected with AdDNSmad2 or AdGFP and treated or untreated with TGFβ1. Relative levels of Scx and Adamtsl2 mRNA were determined by qPCR. Expression was normalized to the housekeeping gene Hprt. Results are shown relative to the untreated control (*) indicates significance, $p < 0.05$, $n = 5$ (K) Cells were infected with AdSmad3 or AdGFP for 48 h and Smad3 expression was verified via immunoblot. (L) Immunoblots were quantified using ImageJ. (*) indicates significance, $p < 0.05$, $n = 3$ (M) RNA was isolated from cells infected with AdSmad3 or AdGFP for 48 h and then treated or untreated with TGFβ1 for 8 h. Relative levels of Scx, Fmod, Adamtsl2, and Prg4 mRNA were determined by qPCR. Results are shown relative to Ad-GFP infected, untreated controls. All mRNA levels were normalized to the housekeeping gene Hprt. (*) indicates significance, $p < 0.05$, $n = 3$. Detailed results from qPCR REST analysis are shown in Tables S4–S6.


Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.



Western Blot


Smad3 is required for TGF- β to regulate expression of Fmod and Adamtsl2. (A) Primary sclerotome was treated with vehicle control or TGF β 1 for 2 h, 8 h or 24 h. Immunoblot was used to determine the levels of pSmad3, pSmad2 and total Smad2/3. α -Tubulin was used as a general loading control. (B) Quantification of pSmad3 and (C) pSmad2 levels relative to total Smad2/3 are shown. Immunoblots were quantified using ImageJ. (*) indicates significance, $p < 0.05$, $n = 3$ for each (D) Sclerotome was pretreated with vehicle control or SIS3 (5 μ M and 10 μ M) for 24 h and then treated with or without TGF β 1 for 8 h. pSmad3 and Smad2/3 protein levels were analyzed via immunoblot. α -Tubulin was used as a general loading control. (E) Smad3 activity was quantified from the immunoblots using ImageJ as the relative levels of pSmad3 normalized to total Smad2/3. (*) indicates significance, $p < 0.05$, $n = 3$ (F) Relative levels of Scx, Fmod, Adamtsl2 and Prg4 mRNA were determined by qPCR after indicated treatment with SIS3. All mRNA levels were normalized to the housekeeping gene Hprt and then analyzed for significance using REST (*). Results are shown relative to TGF β 1 treated cells. $n = 3$ (G) Cells were infected with AdDNSmad2 or AdGFP control virus 48 h before cells were treated with or without TGF β 1 for 8 h. Immunoblot was used to visualize relative levels of pSmad3, pSmad2 and total Smad2/3. α -Tubulin is shown as a general loading control. (H) pSmad3 and (I) pSmad2 activity were quantified using ImageJ as the level of each phosphoSmad normalized to total Smad2/3. (*) indicates significance, $p < 0.05$, $n = 3$ (J) RNA was extracted from cells that were infected with AdDNSmad2 or AdGFP and treated or untreated with TGF β 1. Relative levels of Scx and Adamtsl2 mRNA were determined by qPCR. Expression was normalized to the housekeeping gene Hprt. Results are shown relative to the untreated control (*) indicates significance, $p < 0.05$, $n = 5$ (K) Cells were infected with AdSmad3 or AdGFP for 48 h and Smad3 expression was verified via immunoblot. (L) Immunoblots were quantified using ImageJ. (*) indicates significance, $p < 0.05$, $n = 3$ (M) RNA was isolated from cells infected with AdSmad3 or AdGFP for 48 h and then treated or untreated with TGF β 1 for 8 h. Relative levels of Scx, Fmod, Adamtsl2, and Prg4 mRNA were determined by qPCR. Results are shown relative to Ad-GFP infected, untreated controls. All

mRNA levels were normalized to the housekeeping gene Hprt. (*) indicates significance, $p < 0.05$, $n = 3$. Detailed results from qPCR REST analysis are shown in Tables S4–S6. Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.



Western Blot

TGF- β signaling regulates noncanonical pathways in the sclerotome. Sclerotome was treated with vehicle control or TGF β 1 for 2, 8 or 24 h. Immunoblot was used to determine activity of ERK, p38 and AKT. (B) ERK, (C) P38 and (D) AKT activity was quantified as the relative levels of the phosphoprotein over the total protein as determined from ImageJ scanned blots. α tubulin was used as a general loading control. (*) indicates significance, $p < 0.05$, $n = 3$. Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.

A

Western Blot

AKT and p38 are not required for TGF- β -mediated regulation of fibrous tissue markers. (A) Sclerotome was treated with varying concentrations (1 μ M, 5 μ M, 10 μ M) of P38 inhibitor, BIRB, for 24 h and then treated with TGF β 1 for 8 h.


Immunoblot was used to determine relative levels of the p38 target, p-MAPK-APK2. α -tubulin was used as a loading control. (B) p-MAPK-APK2 levels were quantified using Image J and GraphPad Prism. (*) indicates significance, $p < 0.05$, $n = 3$.

(C) RNA was isolated from cells treated with BIRB and TGF β 1 as indicated. qPCR was used to measure the relative expression of Scx, Fmod, and Adamtsl2. mRNA levels were normalized to the housekeeping gene HPRT. Expression is shown relative to the TGF β 1 treated control. Results were analyzed with REST. (*) indicates significance, $p < 0.05$, $n = 3$.

(D) Sclerotome was treated with varying concentrations (1 μ M, 5 μ M) of AKT inhibitor, MK, for 24 h and then treated with TGF β 1 for 8 h. Immunoblot was used to measure relative levels of pAKT and total AKT. α -tubulin was used as a general control. (E) Immunoblots were scanned using Image J and quantified. Activation of AKT was measured as pAKT over total AKT. (*) indicates significance, $p < 0.05$, $n = 3$


(F) mRNA was isolated from cells treated with MK and TGF β 1 as indicated. qPCR was used to measure the relative levels of Scx, Fmod, or Adamtsl2 mRNA. All mRNA levels were normalized to the housekeeping gene HPRT. Expression is shown relative to the TGF β 1 treated samples. Results were analyzed with REST. (*) indicates significance, $p < 0.05$, $n = 3$.

Detailed results from qPCR REST analysis are shown in Tables S7, S8. Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.




E Western Blot

AKT and p38 are not required for TGF-β-mediated regulation of fibrous tissue markers. (A) Sclerotome was treated with varying concentrations (1 μM, 5 μM, 10 μM) of P38 inhibitor, BIRB, for 24 h and then treated with TGFβ1 for 8 h. Immunoblot was used to determine relative levels of the p38 target, p-MAPK-APK2. α-tubulin was used as a loading control. (B) p-MAPK-APK2 levels were quantified using Image J and GraphPad Prism. (*) indicates significance, $p < 0.05$, $n = 3$. (C) RNA was isolated from cells treated with BIRB and TGFβ1 as indicated. qPCR was used to measure the relative expression of Scx, Fmod, and Adamtsl2. mRNA levels were normalized to the housekeeping gene HPRT. Expression is shown relative to the TGFβ1 treated control. Results were analyzed with REST. (*) indicates significance, $p < 0.05$, $n = 3$. (D) Sclerotome was treated with varying concentrations (1 μM, 5 μM) of AKT inhibitor, MK, for 24 h and then treated with TGFβ1 for 8 h. Immunoblot was used to measure relative levels of pAKT and total AKT. α-tubulin was used as a general control. (E) Immunoblots were scanned using Image J and quantified. Activation of AKT was measured as pAKT over total AKT. (*) indicates significance, $p < 0.05$, $n = 3$ (F) mRNA was isolated from cells treated with MK and TGFβ1 as indicated. qPCR was used to measure the relative levels of Scx, Fmod, or Adamtsl2 mRNA. All mRNA levels were normalized to the housekeeping gene HPRT. Expression is shown relative to the TGFβ1 treated samples. Results were analyzed with REST. (*) indicates significance, $p < 0.05$, $n = 3$. Detailed results from qPCR REST analysis are shown in Tables S7, S8. Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.




Western Blot

ERK is required for fibrous tissue marker regulation but not required to inhibit chondrogenesis. (A) Sclerotome was treated with a MEK inhibitor PD184352, PD, to inhibit ERK activity, for 24 h and then cells were treated with TGFβ1 for 8 h. Immunoblot was used to determine relative levels of pERK1/2, ERK1/2, and α-tubulin. (B) Blots were quantified using ImageJ and Graphpad Prism. Activity of ERK was determined as the ratio of pERK over ERK under the indicated conditions. (*) indicates significance, $p < 0.05$, $n = 3$. (C, D) RNA was isolated from cells under the indicated conditions and qPCR was used to determine the relative expression of (C) Scx, Fmod and Adamtsl2 mRNA or (D) Ebf1 and cMAF mRNA. mRNA levels were normalized to Hprt. Expression is shown relative to the untreated control. Data analyzed by REST software. (*) indicates significance, $p < 0.05$, $n = 3$. Detailed results from qPCR REST analysis are shown in Tables S9, S10. Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.



Western Blot

Scx is required for the expression of Fmod and Adamtsl2. (A) C3H10T1/2 cells were transduced with a scrambled siRNA control (Scram) or Scx specific siRNA for 24 h. Representative immunoblot showing Scx protein. α -tubulin is used as a loading control. (B) Quantification of Scx immunoblots comparing Scram transduced and Scx siRNA transduced cells. Protein levels were normalized to α tubulin and quantified using ImageJ and GraphPad Prism, n = 3, (*) indicates significance, p < 0.05. (C) Scx mRNA levels were determined by qPCR in Scram transduced and Scx siRNA transduced cells. Data was normalized to Hprt mRNA and analyzed using REST software, n = 6, (*) indicates significance, p < 0.05. (D) RNA was collected from cells that were transduced with Scram siRNA or Scx siRNA for 24 h and subsequently either left untreated (-) or treated with TGF β 1(+) for 8 h. Fmod and Adamtsl2 mRNA levels were determined by qPCR. mRNA levels were normalized to Hprt. qPCR data was analyzed by REST software, n = 6, (*) indicates significance, p < 0.05. Detailed results from qPCR REST analysis are shown in Tables S11, S12. Immunoblots were cropped for clarity. Examples of uncropped blots are found in Supplementary Figures. Figure provided by CiteAb. Source: Sci Rep, PMID: 33288795.



Immunofluorescence Microscopy

Immunofluorescence of Mouse Anti-Alpha Tubulin Antibody.

Cell line: HeLa.

Primary Antibody: Alpha Tubulin (p/n 200-301-880) at 4 μ g/mL (1:250) for 1hr at RT.

Secondary Antibody: Goat Anti-Mouse DyLight™ 488 (p/n 610-141-121) at 1 μ g/mL (1:1000) overnight at 4 °C.

Fixative: Ice Cold Methanol.


Permeabilization: Ice Cold Methanol.

Nuclear stain: Hoechst 33342.

Expected Localization: Cytoplasmic.

Image: A) Alpha Tubulin, B) Nuclear Stain, C) Merge, D)

Secondary Only Control.

**Immunofluorescence Microscopy**

Immunofluorescence of Mouse Anti-Alpha Tubulin Antibody.

Cell line: HeLa.

Primary Antibody: Alpha Tubulin (p/n 200-301-880) at 4 $\mu\text{g}/\text{mL}$ (1:250) for 1hr at RT.

Secondary Antibody: Goat Anti-Mouse DyLight™ 488 (p/n 610-141-121) at 0.1 $\mu\text{g}/\text{mL}$ (1:10000) overnight at 4 °C.

Fixative: Ice Cold Methanol.

Permeabilization: Ice Cold Methanol.


Nuclear stain: Hoechst 33342.

Magnification: 40X.

Expected Localization: Cytoplasmic.

Image: HeLa cell nucleus in the anaphase stage of mitosis.

Microtubule-based mitotic spindles are clearly visible.

**Immunofluorescence Microscopy**

Immunofluorescence of Mouse Anti-Alpha Tubulin Antibody.

Cell line: HeLa.

Primary Antibody: Alpha Tubulin (p/n 200-301-880) at 4 $\mu\text{g}/\text{mL}$ (1:250) for 1hr at RT.

Secondary Antibody: Goat Anti-Mouse DyLight™ 488 (p/n 610-141-121) at 0.1 $\mu\text{g}/\text{mL}$ (1:10000) overnight at 4 °C.

Fixative: Ice Cold Methanol.

Permeabilization: Ice Cold Methanol.


Nuclear stain: Hoechst 33342.

Magnification: 40X.

Expected Localization: Cytoplasmic.

Image: HeLa cell nucleus in the metaphase stage of mitosis.

Microtubule-based mitotic spindles are clearly visible.



Immunofluorescence Microscopy

Immunofluorescence of Mouse Anti-Alpha Tubulin Antibody.

Cell line: HeLa.

Primary Antibody: Alpha Tubulin (p/n 200-301-880) at 4 $\mu\text{g}/\text{mL}$ (1:250) for 1hr at RT.

Secondary Antibody: Rabbit Anti-Mouse Texas Red™ (p/n 610-409-C49) at 1 $\mu\text{g}/\text{mL}$ (1:1000) overnight at 4 °C.


Fixative: Ice Cold Methanol.

Permeabilization: Ice Cold Methanol.

Nuclear stain: Hoechst 33342.

Expected Localization: Cytoplasmic.

Image: A) Alpha Tubulin, B) Nuclear Stain, C) Merge, D) Secondary Only Control.



Immunofluorescence Microscopy

Rockland's α -tubulin monoclonal antibody detects tubulin (colored RED) in STED immunofluorescence microscopy.

Methanol fixed A431 cells were blocked with normal goat serum. The cells were then probed with 0.4 $\mu\text{g}/\text{mL}$ final

concentration of anti- α -tubulin and detected with 0.2 $\mu\text{g}/\text{mL}$ ATTO 425 conjugated anti-Mouse IgG [GOAT] (p/n 610-151-121) secondary antibody. Also shown in this 2-color STED

image is Rockland's Anti-HDAC-1 [RABBIT] (p/n 600-401-879) detected with DyLight™488 conjugated Anti-RABBIT IgG [GOAT] secondary antibody (colored GREEN). Image

courtesy of Myriam Gastard, Leica Microsystems, USA.

Western Blot

HeLa whole cell lysate (p/n W09-000-364) [left lane] and HEK293 whole cell lysate (W09-000-365) [right lane] were


loaded with 10 μg of lysate each. The blot was blocked with Blocking Buffer (p/n MB-070) for 30 min at RT, then washed


and incubated with Rockland's anti-Tubulin monoclonal antibody diluted in Blocking Buffer (p/n MB-070) at 1:1,000

for 1 h at RT. After washing, blot was incubated with a 1:40,000 dilution of Rockland's HRP Rb a-Ms IgG (p/n 610-4302) secondary antibody in Blocking Buffer (p/n MB-070)


for 30 minutes at RT. Data was collected using Bio-Rad

VersaDoc® 4000 MP.




**Immunofluorescence Microscopy**


Anti-alpha-Tubulin (MOUSE) Monoclonal Antibody.
Immunofluorescence image showing MCF-7 cell staining of Anti-alpha-Tubulin (MOUSE) Monoclonal Antibody - (p/n 200-301-880) in green and staining of Anti-Gli-3 (RABBIT) Antibody - (p/n 600-401-694) in red.

**Immunofluorescence Microscopy**

Anti-alpha-Tubulin (MOUSE) Monoclonal Antibody.
Immunofluorescence image one showing MCF-7 cell staining of Anti-alpha-Tubulin (MOUSE) Monoclonal Antibody - (p/n 200-301-880) in green. Image two showing MCF-7 cell staining of Anti-Gli-3 (RABBIT) Antibody - (p/n 600-401-694) in red. Image three showing MCF-7 cell superimposed staining of Anti-alpha-Tubulin (MOUSE) Monoclonal Antibody - (p/n 200-301-880) in green and staining of Anti-Gli-3 (RABBIT) Antibody - (p/n 600-401-694) in red.

**Immunofluorescence Microscopy**


Rockland's anti- α -tubulin monoclonal antibody was used at a 0.1 μ g/mL to detect tubulin in 4% paraformaldehyde fixed A459 cells. Staining is shown using conventional confocal microscopy (left panel) and by high resolution TCS STED nanoscopy (right panel). DyLight488™ conjugated anti-mouse IgG secondary antibody (p/n 610-141-121) was used for detection at 1 μ g/mL. Personal Communication, Myriam Gastard, Leica Microsystems, Exton PA.


**Immunofluorescence Microscopy**

Immunofluorescence microscopy of Anti-tubulin in U-87 MG cells using FITC-conjugated Fluorescent TrueBlot® anti-mouse IgG (p/n 18-0217-32) for detection. U87-MG cells were fixed with 100% methanol, permeabilized with 0.3% Triton X-100, blocked with 5% rat serum/0.3 Triton X-100 for 1 hr, then incubated with 15 µg/mL of anti- α -tubulin primary antibody (p/n 200-301-880) at 4°C overnight. Following 3 washes in 1X PBS for 5min each, 5µg/mL of Fluorescent TrueBlot® anti-mouse IgG Fluorescein (p/n 18-0217-32) was added and allowed to incubate for 1hr at room temperature. 5 µg/mL of Fluorescent TrueBlot® anti-mouse IgG FITC was added and allowed to incubate for 1hr at room temperature. Nucleus was counterstained with DAPI present in mounting medium. The predicted main localization is microtubules. Image taken at 63X magnification. Merged α -tubulin (green)/DAPI (blue) image shown.

Immunofluorescence Microscopy


Immunofluorescence microscopy of α -tubulin in U-87 MG cells using FITC-conjugated Fluorescent TrueBlot® anti-mouse IgG (p/n 18-0217-32) for detection. U87-MG cells were fixed with 100% methanol, blocked (5% rat serum/0.3% Triton X-100) for 1hr, then incubated with 15µg/mL of anti- α -tubulin primary antibody (p/n 200-301-880) at 4°C overnight. Following 3 washes in 1X PBS for 5min each, 5µg/mL of Fluorescent TrueBlot® anti-mouse IgG Fluorescein was added and allowed to incubate for 1hr at room temperature. 5µg/mL of Fluorescent TrueBlot® anti-mouse IgG FITC was added and allowed to incubate for 1hr at room temperature. Nucleus was counterstained with DAPI present in mounting medium. The predicted main localization is microtubules. Image taken at 63X magnification. (a) Merged α -tubulin (green)/DAPI (blue) image shown (b) secondary only.






Immunofluorescence Microscopy

Immunofluorescence microscopy of α -tubulin in HeLa cells using DyLight™ 680-conjugated Fluorescent TrueBlot® anti-mouse IgG (p/n 18-4417-32) for detection. HeLa cells were fixed with 100% methanol, blocked (5% rat serum/0.3% Triton X-100 in 1X PBS) for 1hr, then incubated with 15 μ g/mL of anti-alpha-tubulin primary antibody (p/n 200-301-880) at 4°C overnight. Following 3 washes in 1X PBS for 5 min each, 5 μ g/mL of Fluorescent TrueBlot® anti-mouse IgG DyLight™ 680 was added and allowed to incubate for 1hr at room temperature. Nuclei were counterstained with DAPI present in mounting medium. The predicted main localization is microtubules. Image taken at 63X magnification. (a) Merged α -tubulin (red)/DAPI (blue) image shown. (b) secondary antibody only.




Immunofluorescence Microscopy

Rockland mouse monoclonal anti-tubulin antibody (p/n 200-301-880) was used with ATTO 425 Goat anti-mouse (p/n 610-151-121, shown in red) to detect tubulin by immunofluorescence. DyLight 488 Goat anti-Rabbit (p/n 611-141-122, shown in green) was used in the same experiment to detect rabbit anti-Clathrin polyclonal antibody. Data was collected on a STED-CW TCS-SP5 Confocal system (Leica Microsystems) equipped with a DFC 350FX Camera allowing sequential acquisition in wide-field, confocal and STED CW imaging modes and provided courtesy of: Myriam Gastard, PhD, personal communication, Leica Microsystems, Inc. USA




Immunohistochemistry

Rockland's anti- α -tubulin monoclonal antibody was used at a 2.5 μ g/mL to detect tubulin in thyroid follicular epithelium (40X) showing moderate to strong cytoplasmic staining (image). Moderate to strong cytoplasmic staining was also observed within subsets of neurons and glia, and epithelial cells including adrenal, breast, colon, pancreas, kidney, prostate, placenta, skin, testis, uterus, thyroid, and within lymphoid organs. The image shows the localization of the antibody as the precipitated red signal, with a hematoxylin purple nuclear counterstain. Tissue was formalin-fixed and paraffin embedded. Personal Communication, Vasiliki Demas, LifeSpan Biosciences, Seattle, WA.




Immunohistochemistry

Immunohistochemistry of Mouse anti-Alpha-Tubulin antibody. Tissue: human colon. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: anti-Alpha-Tubulin antibody at 5 µg/mL for 1 h at RT. Secondary antibody: Peroxidase mouse secondary antibody at 1:10,000 for 45 min at RT. Staining: Alpha-Tubulin as precipitated red signal with hematoxylin purple nuclear counterstain.



Immunofluorescence Microscopy

Immunofluorescences of Mouse monoclonal anti-Alpha Tubulin antibody. Cell Type: A431 cells. Fixation: 4% paraformaldehyde 10 min. Permeabilization: 0.5% Triton X 30 min. Primary Ab: (p/n 200-301-880, lot 28977) 1:250 X 30 min. Secondary Ab: (p/n 610-142-121, lot 21290) 1:1000 Overnight 4°C.



Immunofluorescence Microscopy

Immunofluorescence of Mouse monoclonal anti-Alpha Tubulin antibody. Cell Type: A431 cells. Fixation: 4% paraformaldehyde 10 min. Permeabilization: 0.5% Triton X 30 min. Primary Ab: (p/n 200-301-880, lot 28977) 1:250 72 hours 4°C. Secondary Ab: Gt anti-Mouse DyLight™488 (p/n 610-141-121, lot 21286) 1:1000 overnight 4°C.

References

- de la Calle-Fabregat C et al. NF- κ B and TET2 promote macrophage reprogramming in hypoxia that overrides the immunosuppressive effects of the tumor microenvironment. *Sci Adv.* (2024)
- Hoffmeister H et al. Characterization of the nuclear import of the human CHD4-NuRD complex. *J Cell Sci.* (2023)
- Alburquerque-Bejar JJ et al. MYC activation impairs cell-intrinsic IFN γ signaling and confers resistance to anti-PD1/PD-L1 therapy in lung cancer. *Cell Rep Med.* (2023)
- Clayton, SW et al. TGF β signaling is required for sclerotome resegmentation during development of the spinal column in Gallus gallus. *Developmental Biology* (2022)
- Diggins NL et al. Human Cytomegalovirus miR-US25-1 Targets the GTPase RhoA To Inhibit CD34+ Hematopoietic Progenitor Cell Proliferation To Maintain the Latent Viral Genome. *mBio.* (2021)
- Clayton SW et al. TGF β signaling is required for sclerotome resegmentation during development of the spinal column in Gallus gallus. *bioRxiv Preprint* (2021)
- Diggins NL et al. A Novel miRNA-Mediated Mechanism to Retain Latent HCMV Genomes in Hematopoietic Progenitor Cells. *Cell Press Preprint* (2021)
- Wu G et al. A two-layered targeting mechanism underlies nuclear RNA sorting by the human exosome. *Cell Rep.* (2020)
- Wu M et al. The RNA exosome shapes the expression of key protein-coding genes. *Nucleic Acids Res.* (2020)
- Clayton SW et al. Canonical and noncanonical TGF- β signaling regulate fibrous tissue differentiation in the axial skeleton. *Sci Rep.* (2020)
- Nguyen T et al. Uncovering the role of N-acetyl-aspartyl-glutamate as a glutamate reservoir in cancer. *Cell Rep.* (2019)
- Roy K et al. Polycystin-1, the product of the polycystic kidney disease gene PKD1, is post-translationally modified by palmitoylation. *Mol Biol Rep.* (2018)
- Xie et al. Cardiac glycoside bufalin blocks cancer cell growth by inhibition of Aurora A and Aurora B activation via PI3K-Akt pathway. *Oncotarget* (2018)
- Chavez et al. SOX9 protein is stabilized by TGF- β and regulates PAPSS2 mRNA expression in chondrocytes. *Osteoarthritis Cartilage* (2017)
- Patel et al. The Common R71H-G230A-R293Q Human TMEM173 Is a Null Allele. *Journal of Immunology* (2017)
- Roy et al. Palmitoylation of the ciliary GTPase ARL13b is necessary for its stability and its role in cilia formation. *Journal of Biological Chemistry* (2017)
- Laitem et al. CDK9 inhibitors define elongation checkpoints at both ends of RNA polymerase II-transcribed genes. *Nature Structural & Molecular Biology* (2015)
- Reisen F et al. Linking phenotypes and modes of action through high-content screen fingerprints. *Assay Drug Dev Technol.* (2015)
- Blaauboer SM et al. MPYS/STING-mediated TNF- α , not type I IFN, is essential for the mucosal adjuvant activity of (3'-5')-cyclic-di-guanosine-monophosphate in vivo. *J Immunol.* (2014)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.