

Datasheet for 200-301-A87**Mesothelin Antibody****Overview**

Description:	Anti-Mesothelin (MOUSE) Monoclonal Antibody - 200-301-A87
Item No.:	200-301-A87
Size:	100 µg
Applications:	FC, IHC, WB
Reactivity:	Human
Host Species:	Mouse

Product Details

Background:	Anti Mesothelin Antibody recognizes Mesothelin that is a glycosyl-phosphatidylinositol–anchored glycoprotein present on the cell surface of various human solid tumors. The mesothelin (MSLN) gene encodes a 71-kDa precursor protein that is processed to a 40-kDa glycosylphosphatidylinositol–anchored protein that composes the mature portion and an NH2 terminal 31-kDa fragment called megakaryocyte-potentiating factor that is released from the cell. Mesothelin is a tumor differentiation antigen present at low levels on a restricted set of normal adult tissues, such as mesothelium, but aberrantly over expressed in mesotheliomas, ovarian, and pancreatic cancers. The biological functions of mesothelin remain elusive. A recent study showed that mesothelin binds to MUC16/CA125, and that this interaction mediates cell adhesion, suggesting that there may be an important role for MUC16/CA125 and mesothelin in the metastatic spread of ovarian cancer.
Synonyms:	mouse anti-Mesothelin Antibody, Mesothelian, MN, MB, Pre-pro-megakaryocyte-potentiating factor, CAK1 antigen
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	MB-G10
Format:	IgG2a

Target Details

Gene Name:	MSLN
-------------------	------

Reactivity:	Human
Immunogen Type:	Recombinant Protein
Immunogen:	This antibody was produced in mesothelin-deficient mice by immunizations with plasmid cDNA encoding human MSLN full length protein followed by a single boost of a recombinant human mesothelin-Fc fusion protein.
Purity/Specificity:	This antibody is directed against human mesothelin protein. This product was purified from tissue culture supernatant fluid by Protein A chromatography. Cross reactivity with homologues from other sources has not been tested.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - Q13421• NCBI - 53988378• GeneID - 10232

Application Details

Tested Applications:	FC, IHC, WB
Application Note:	This antibody has been tested for use in immunohistochemistry and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 40 kDa in size corresponding to mature mesothelin by western blotting in the appropriate cell lysate or extract. For Anti-mesothelin immunohistochemistry, archival PEFF human tissues were deparaffinized followed by hydration. Antigen-retrieval is recommended. Block tissues with 1% BSA in PBS for 30 min at 23° C. Antibodies are diluted in 1% BSA and reacted with tissue for 60 min at room temperature.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:50,000
FC:	1:200
IHC:	1:100
WB:	1:1,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store antibody at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images




Flow Cytometry

Flow Cytometry Results of Anti-Mesothelin (MOUSE) Monoclonal Antibody. The green histogram shows NCI-H226 cells and blue histogram shows MCF-7 cells. Both cell lines are stained with a 1:200 dilution Anti-Mesothelin (MOUSE) Monoclonal Antibody. The secondary antibody use was Anti-Mouse IgG (H&L) (GOAT) Antibody DyLight™ 488 Conjugated (p/n 610-141-002, lot#43322) at the 1:400 dilution.




Flow Cytometry

Cells were incubated with the indicated primary antibodies or isotope control antibodies, and appropriate secondary antibody (goat anti-mouse IgG, R-PE or goat anti-mouse, FITC). Results are shown as histogram plots for binding of primary antibody (black trace) or isotype control (gray trace). A431 cells are negative for both MSLN and CA125. A549 cells show low mesothelin expression (3.5×10^3 sites/cell) but highly express CA125. Fig3. PMID: 25118887




Flow Cytometry

(A) Tumor cells present in ascites were evaluated by their positivity for the tumor differentiation antigen mesothelin by flow cytometry using mouse anti-mesothelin (p/n 200-301-A87) primary antibody followed by goat anti-mouse IgG PE labeled secondary antibody. Shaded histogram represents the binding of isotype control antibody and solid histograms represent the binding of anti-mesothelin antibody. (B) Reactivity of lymphocytes with autologous tumor cells was evaluated by measuring IFN-γ released in the supernatants of co-cultures of lymphocytes and tumor cells. (C) Flow cytometry analysis of PD-L1 expression on autologous tumor cells obtained from ascites of patient NCI-Meso29 grown in the presence (solid) or absence (shaded) of autologous lymphocytes cultured from ascites and 106 IU/mL IL-2. Fig 4. PMID: 27544053




Immunohistochemistry

MSLN immunohistochemistry in colon cancer tissue. (A) Case of tubular adenocarcinoma. Diffuse and luminal MSLN expression was detected. (B) Case of poorly differentiated adenocarcinoma exhibiting solid/sheet-like proliferation with undetectable MSLN expression. Left, low-power magnification; right, high-power magnification. Scale bar, 500 μm. MSLN, mesothelin. Fig 1. PMID: 32194667




Western Blot

Expression of MSLN in cultured colon cancer cells. MSLN was expressed in four of seven colon cancer cell lines with no association with sex. MSLN, mesothelin; F, female; M, male. Fig 3. PMID: 32194667




Immunohistochemistry

Immunohistochemistry using Rockland's anti-mesothelin antibody to react with two epitopes on mesothelin in PEFF human mesothelioma tissue sections treated by antigen retrieval methods. Anti-mesothelin primary antibodies were used at 10 µg/mL to label these sections as follows: C, MAb MB; and D, MAb MN followed by goat anti-mouse IgG conjugated to horseradish peroxidase at 25 µg/mL in 1% BSA/PBS for 30 minutes. (magnification, ×200; bar, 50 µm). Reprinted with permission from Clin.Cancer Res. 11(16):5840-6.



SDS-PAGE

SDS-PAGE of Mouse anti-Mesothelin Monoclonal Antibody. Lane 1: Non-Reduced Mouse anti-Mesothelin Monoclonal Antibody. Lane M: 3 µL OPAL Pre-stained Marker (p/n MB-210-0500). Lane 2: Reduced Mouse anti-Mesothelin Monoclonal Antibody. Load: 1 µg per lane. Predicted/Observed size: Non-reduced at 160 kDa; Reduced at 55, 25 kDa.



Western Blot

Western blotting using Rockland's anti-mesothelin antibody. Load: Mesothelin-Fc (lane 1, 100 ng; lane 2, 25 ng; lane 3, 6 ng; lane 4, 2 ng; and lane 5, 0.4 ng) and CD25-Fc (lane 6, 50 ng) Primary antibody: anti-mesothelin at 1mg/ml. Secondary Antibody: ALP goat anti-mouse IgG and BCIP/NBT substrate. Reprinted with permission from Clin.Cancer Res. 11(16):5840-6.

References

- Koukoulis K, et al. Targeting Mesothelin via the native T cell receptor. *BioRxiv (Preprint)* (2025)
- Marin-Muller C et al. Nanoparticle-Mediated Therapy with miR-198 Sensitizes Pancreatic Cancer to Gemcitabine Treatment through Downregulation of VCP-Mediated Autophagy. *Pharmaceutics*. (2023)
- Inoue S. et al. Diffuse mesothelin expression leads to worse prognosis through enhanced cellular proliferation in colorectal cancer. *Oncology Letters* (2020)
- Leshem, Y et al. Combining Local Immunotoxins Targeting Mesothelin with CTLA-4 Blockade Synergistically Eradicates Murine Cancer by Promoting Anticancer Immunity. *Cancer Immunology Research* (2017)
- Illei, PB et al. Mesothelin Expression in Advanced Gastroesophageal Cancer Represents a Novel Target for Immunotherapy. *Applied Immunohistochemistry & Molecular Morphology : Aimm* (2016)
- Awuah et al. Reduced Shedding of Surface Mesothelin Improves Efficacy of Mesothelin-Targeting Recombinant Immunotoxins. *Molecular Cancer Therapeutics* (2016)
- Khanna et al. Malignant Mesothelioma Effusions Are Infiltrated by CD3+ T Cells Highly Expressing PD-L1 and the PD-L1+ Tumor Cells within These Effusions Are Susceptible to ADCC by the Anti-PD-L1 Antibody Avelumab. *Journal of Thoracic Oncology* (2016)
- Alewine, C et al. Efficacy of RG7787, a next-generation mesothelin-targeted immunotoxin, against triple-negative breast and gastric cancers. *Molecular Cancer Therapeutics* (2014)
- Zhang, J et al. Megakaryocytic potentiating factor and mature mesothelin stimulate the growth of a lung cancer cell line in the peritoneal cavity of mice. *PLoS One* (2014)
- Zhang, J et al. Loss of mesothelin expression by mesothelioma cells grown in vitro determines sensitivity to anti-mesothelin immunotoxin SS1P. *Anticancer Research* (2012)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.