

Datasheet for 200-301-BX1**IL-33 Antibody [12H6D12]****Overview**

Description:	Anti-IL-33 (MOUSE) Antibody - 200-301-BX1
Item No.:	200-301-BX1
Size:	100 µg
Applications:	ELISA, IHC, WB
Reactivity:	H. sapiens (Human), Mus musculus (Mouse)
Host Species:	Mouse

Product Details

Background: Interleukin-33 (IL-33) is a recently identified member of the IL-1 family of cytokines whose other members include IL-1a/beta, IL-1Ra and IL-18. Its receptor has been shown to be ST2, an IL-1 receptor family member that also acts as a negative regulator of TLR-IL-1R signaling and IL-1R accessory protein (IL-1RAcP). Receptor binding of IL-33 activates NF-kB and MAP kinases and induces the expression of TH2-associated cytokines such as IL-4, IL-5 and IL-6. Prolonged IL-33 treatment of mice led to the development of eosinophilia, splenomegaly, and severe pathological changes in mucosal organs such as lungs, esophagus and small intestine. Recent experiments have shown that IL-33 can also co-localize with heterochromatin and possesses transcriptional repressor activities, indicating that IL-33 may function as both a proinflammatory cytokine and an intracellular nuclear factor with transcriptional regulatory properties.

Synonyms:	IL-33 Antibody [12H6D12] , DVS27, IL1F11, NF-HEV, NFEHEV, C9orf26
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	[12H6D12]
Format:	IgG1

Target Details

Gene Name:	IL33
Reactivity:	H. sapiens (Human), Mus musculus (Mouse)

Immunogen Type:	Recombinant Protein
Immunogen:	Anti-IL-33 antibody was produced in mice by repeated immunizations with the recombinant IL-33 protein.
Purity/Specificity:	Anti-IL-33 Monoclonal Antibody was Protein A purified from monospecific antiserum by immunoaffinity purification. Cross reactivity with IL-33 [12H6D12] from other sources has not been determined.
Relevant Links:	• UniProtKB - O95760 • GeneID - 90865 • NCBI - NP_254274

Application Details

Tested Applications:	ELISA, IHC, WB
Application Note:	Anti-IL-33 Antibody has been tested for use in ELISA, Western Blotting, and Immunohistochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 31 kDa in Western Blots of specific cell lysates and tissues. Validated in human samples. All other applications and species not yet tested.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000
IHC:	5 µg/mL
WB:	1 µg/mL

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/ml by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.


Expiration: Expiration date is one (1) year from date of receipt.

Images



Immunohistochemistry

Immunohistochemistry of Mouse anti-IL-33 antibody. Tissue: human lymph node. Primary antibody: IL-33 antibody at 5 µg/mL. Secondary antibody: Peroxidase mouse secondary antibody at 1:5,000. Localization: IL-33 is secreted and nuclear.



Western Blot

Western Blot of Mouse anti-IL-33 antibody. Lane A: 125ng of recombinant IL-33. Primary antibody: IL-33 antibody at 1 µg/mL overnight at 4°C. Secondary antibody: Mouse HRP secondary antibody. Block: 5% BLOTTO. Predicted/Observed size: 29 kDa, 29 kDa for IL-33.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.