

Datasheet for 200-301-E63**BIN1 Antibody****Overview**

Description:	Anti-BIN1 (MOUSE) Monoclonal Antibody - 200-301-E63
Item No.:	200-301-E63
Size:	100 µg
Applications:	ELISA, IHC, WB
Reactivity:	Mouse
Host Species:	Mouse

Product Details

Background: Bin1 is a conserved member of the BAR family of genes that have been implicated in diverse cellular processes including endocytosis, actin organization, programmed cell death, stress responses, and transcriptional control. The first mammalian BAR protein to be discovered, Amphiphysin I (AmphI), was identified in an immunoscreen for proteins associated with the plasma membranes of synaptic neurons, functions in the control of clathrin-dependent synaptic vesicle endocytosis. The mammalian Bin1 gene was first identified in a two hybrid screen for polypeptides that bind to the N-terminal Myc box 1 (MB1) portion of the c-Myc oncoprotein. Bin1 is similar to AmphI in overall structure, with an N-terminal BAR domain and a C-terminal SH3 domain. However, the Bin1 gene is more complex than the AmphI gene, encoding at least seven different splice variants that differ widely in subcellular localization, tissue distribution, and ascribed functions. Alternate splicing of the Bin1 gene results in ten transcript variants encoding different isoform. Bin1 is expressed ubiquitously in mammalian cells. Certain splice variants of Bin1 are expressed in the neurons, muscle cells or tumor cells.) Bin1 may act with cancer suppressor and inhibits malignant cell transformation. Studies in mouse suggest that this gene plays an important role in cardiac muscle development. Bin1 has also been implicated in Alzheimer disease and cardiac disease. Defects in BIN1 are the cause of centronuclear myopathy autosomal recessive; also known as autosomal recessive myotubular myopathy.

Synonyms:	mouse anti-BIN1 Antibody, AMPHL, Myc box-dependent-interacting protein 1, Amphiphysin II, Amphiphysin-like protein, Box-dependent myc-interacting protein 1, Bridging integrator 1, BIN 1, BIN-1, BIN1 antibody, anti-BIN1 antibody
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	2F11

Format: IgG1

Target Details

Gene Name: BIN1

Reactivity: Mouse

Immunogen Type: Recombinant Protein

Immunogen: Anti-BIN1 (MOUSE) Monoclonal Antibody was produced in mouse by repeated immunizations with chimeric protein that encoded the human BIN1 BAR domain followed by hybridoma development.

Purity/Specificity: Anti-BIN1 was purified from clarified mouse ascetic fluid by Protein A chromatography followed by extensive dialysis against the buffer stated above. BIN1 antibody is specific for human BIN1 protein. A BLAST analysis was used to suggest cross-reactivity with BIN1 from human and mouse sources based on 100% homology with the immunizing sequence. Cross-reactivity with BIN1 from other sources has not been determined.

Relevant Links:

- [GenelD - 274](#)
- [NCBI - NP_004296.1](#)
- [UniProtKB - O00499](#)

Application Details

Tested Applications: ELISA, IHC, WB

Application Note: Anti-BIN1 antibody has been tested for use in Western Blot, ELISA, IP, and IHC. This antibody is suitable for. Specific conditions for reactivity should be optimized by the end user.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:5000-1:50000

IHC: 1:100-1:500

IP: 10-100 μ L

WB: 1:500-1:1500

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: 0.01% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images




Western Blot

Immunoblot analysis of T-tubule and SR proteins. a Representative immunoblots of microsomal fractions from LV tissues of each study group that were probed with anti-BIN1, Jph2, Cav1.2 and RyR2 antibodies. Observed molecular weights (kDa) of each protein are indicated. Ponceau red staining of the immunoblots are shown with molecular weight markers. b,c Bar graphs show the quantitation of the luminescence intensity of the indicated protein bands, normalized to Ponceau red staining and expressed relative to the mean sham values. Values are mean \pm SD for n = 4–6 hearts/group. Statistical analysis using one-way ANOVA with post-hoc Tukey's multiple group comparisons; p-values are indicated.


Fig 6.

PMID: 31810440



Immunohistochemistry

Immunohistochemistry of Mouse Anti-BIN1 antibody.
 Tissue: skeletal muscle from BIN-1 wild type (+/+) and null (-/-) mouse. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: BIN-1 antibody for 1hr at RT. Secondary antibody: Peroxidase mouse secondary antibody at 1:10,000 for 45 min at RT. Localization: BIN1 is nuclear and cytoplasmic. Staining: BIN1 as precipitated brown signal.



Western Blot

Western Blot of Mouse Anti-BIN-1 antibody. Lane 1: Keratinocyte derived from Bin-1 wild type. Lane 2: Keratinocyte derived from Bin-1 null mice. Load: 35 µg per lane. Primary antibody: BIN-1 antibody at 1:400 for overnight at 4°C. Secondary antibody: IRDye800™ mouse secondary antibody at 1:10,000 for 45 min at RT. Block: 1xPBS, 0.4% Tween-20 (PBS/T) overnight at 4°C. Bin1 isoforms (+exon13, -exon13). *indicates non-specific signal.

References

- An S et al. Adverse transverse-tubule remodeling in a rat model of heart failure is attenuated with low-dose triiodothyronine treatment. *Mol Med.* (2019)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.