

Datasheet for 200-301-F60**HSP60 Antibody****Overview**

Description:	Anti-HSP60 (MOUSE) Monoclonal Antibody - 200-301-F60
Item No.:	200-301-F60
Size:	200 µg
Applications:	FC, IHC, IP, WB
Reactivity:	Human, Mouse, Rat, Soybean
Host Species:	Mouse

Product Details**Background:**

In both prokaryotic and eukaryotic cells, the misfolding and aggregation of proteins during biogenesis and under conditions of cellular stress are prevented by molecular chaperones. Members of the HSP60 family of heat shock proteins are some of the best characterized chaperones. Hsp60, also known as Cpn60 or GroEl, is an abundant protein synthesized constitutively in the cell that is induced to a higher concentration after brief cell shock. It is present in many species and exhibits a remarkable sequence homology among various counterparts in bacteria, plants, and mammals with more than half of the residues identical between bacterial and mammalian Hsp60. Whereas mammalian Hsp60 is localized within the mitochondria, plant Hsp60, or otherwise known as Rubisco-binding protein, is located in plant chloroplasts.

It has been indicated that these proteins carry out a very important biological function due to the fact that Hsp60 is present in so many different species. The common characteristics of the Hsp60s from the divergent species are i) high abundance, ii) induction with environmental stress such as heat shock, iii) homo-oligomeric structures of either 7 or 14 subunits which reversibly dissociate in the presence of Mg²⁺ and ATP, iv) ATPase activity and v) a role in folding and assembly of oligomeric protein structures. These similarities are supported by recent studies where the single-ring human mitochondrial homolog, Hsp60 with its co-chaperonin, Hsp10 were expressed in a *E. coli* strain, engineered so that the groE operon is under strict regulatory control. This study has demonstrated that expression of Hsp60-Hsp10 was able to carry out all essential in vivo functions of GroEL and its co-chaperonin, GroES. Another important function of Hsp60 and Hsp10 is their protective functions against infection and cellular stress. Hsp60 has however been linked to a number of autoimmune diseases, as well as Alzheimer's, coronary artery diseases, MS, and diabetes. Anti-Hsp60 Antibody is useful for researchers interested in Heat Shock research.

Synonyms: 60 kDa heat shock protein, mitochondrial, 60 kDa chaperonin, Chaperonin 60, CPN60, Heat shock protein 60, HSP-60, Hsp60, Mitochondrial matrix protein P1, P60 lymphocyte protein, PD1, CPN60 Antibody, GROEL Antibody, HLD4 Antibody, HSP 60 Antibody, HSP65 Antibody, HSPD1 Antibody, HuCHA60 Antibody, SPG 13 Antibody

Host Species: Mouse

Clonality: Monoclonal

Clone ID: LK1

Format: IgG1

Target Details

Gene Name: HSPD1

Reactivity: Human, Mouse, Rat, Soybean

Immunogen Type: Recombinant Protein

Immunogen: HSP60 Antibody was produced in mice by repeated immunizations raised against recombinant human Hsp60.

Purity/Specificity: Anti-HSP60 Antibody was purified by Protein G chromatography. A BLAST analysis was used to suggest cross-reactivity with Hsp60 from Human, Mouse, Rat, Bovine, Canine, Chicken, Drosophila, Guinea Pig, Hamster, Monkey, Pig, Rabbit, Sheep, Plant, Silkworm, and Xenopus sources based on 100% homology with the immunizing sequence. Cross-reactivity with Hsp60 from other sources has not been determined.

Relevant Links:

- [NCBI - NP_002147.2](#)
- [GeneID - 3329](#)
- [UniProtKB - P10809](#)

Application Details

Tested Applications: FC, IHC, IP, WB

Application Note: Anti-Hsp60 Antibody is tested for WB, IP, ELISA, Flow Cytometry, and IHC-P. Expect a band approximately ~60kDa protein corresponding to the molecular mass of Hsp60 on SDS PAGE immunoblots. Specific conditions for reactivity should be optimized by the end user.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:200

FC: User Optimized

IF:	1:100
IHC:	1:100
IP:	1:200
WB:	1:20000


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.09% (w/v) Sodium Azide
Stabilizer:	50% (v/v) Glycerol

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images




Immunohistochemistry

Immunohistochemistry of mouse anti-Hsp60 antibody.

Tissue: human skin fibroblasts. Antigen retrieval: Left: control, Right: 24 hours after 7th passage of senescence.

Primary Antibody: Hsp60 antibody at 1ug/ml for 1h at RT.

Secondary antibody: Peroxidase mouse secondary at 1:10,000 for 45 min at RT. Localization: Nuclear. Staining: Hsp60 as brown signal.



Western Blot

Western Blot of mouse anti-Hsp60 antibody. Lane 1: A431. Lane 2: A549. Lane 3: HCT116. Lane 4: HeLa. Lane 5: HEK293. Lane 6: HepG2. Lane 7: HL-60. Lane 8: HUVEC. Lane 9: Jurkat. Lane 10: MCF7. Lane 11: PC3. Lane 12: T98G. Lane 13: Rat Brain. Load: 10 µg per lane. Primary antibody: Hsp60 antibody at 1:1000 for overnight at 4°C. Secondary antibody: IRDye800™ mouse secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 61.1 kDa for Hsp60. Other band(s): none.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.