

Datasheet for 200-301-G37**Rhodopsin Antibody****Overview**

Description:	Anti-Rhodopsin (MOUSE) Monoclonal Antibody - 200-301-G37
Item No.:	200-301-G37
Size:	100 µg
Applications:	IF, IHC, IP, WB
Reactivity:	Human, Rat
Host Species:	Mouse

Product Details

Background:	Rhodopsin consists of the protein moiety opsin and a reversibly covalently bound cofactor, retinal. Opsin, a bundle of seven membrane embedded alpha-helices, binds retinal, a photoreactive chromophore, in a central pocket. In addition to being the pigment of the retina that is responsible for both the formation of the photoreceptor cells, its function is to specifically convey information stored in the specific geometry of the chromophore to the surface of the molecule upon light absorption. In the active state, rhodopsin activates transduction, a GTP binding protein. Once activated, transduction promotes the hydrolysis of cGMP by phosphodiesterase. Rhodopsin's activity is believed to be shut off by its phosphorylation followed by binding of the soluble protein arrestin. Mutations in the rhodopsin gene lead to retinitis pigmentosa, which can be inherited as an autosomal dominant, an autosomal recessive or an X-linked recessive disorder. Signaling and GTPase research.
Synonyms:	OPN2 Antibody, opsd Antibody, opsin 2 Antibody, opsin 2 rod pigment Antibody, opsin2 Antibody, RHO Antibody, RP4 Antibody, MGC138309 Antibody, Retinitis Pigmentosa 4 Antibody, Rhodopsin Antibody
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	1D4
Format:	IgG1

Target Details

Gene Name:	RHO
-------------------	-----

Reactivity:	Human, Rat
Immunogen Type:	Other
Immunogen:	Rhodopsin Antibody was produced in mice by repeated immunizations raised against bovine Rhodopsin.
Purity/Specificity:	Anti-Rhodopsin Antibody was purified by Protein G chromatography. A BLAST analysis was used to suggest cross-reactivity with Rhodopsin from all vertebrates based on 100% homology with the immunizing sequence. Cross-reactivity with Rhodopsin from other sources has not been determined. No known reactivity to other proteins. Recognizes Rhodopsin (native and recombinant forms). Binds specifically to the C-terminal epitope -T-E-T-S-Q-V-A-P-A-(COOH).
Relevant Links:	• NCBI - NP_001014890.1 • GeneID - 509933 • UniProtKB - P02699

Application Details

Tested Applications:	IF, IHC, IP, WB
Application Note:	Anti-Rhodopsin Antibody is tested for use in IF microscopy, IHC-P, IP, WB, and ELISA. Expect a band approximately ~40kDa corresponding to specific lysates. Binds specifically to the N-terminus of Rhodopsin. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:200
IF:	User Optimized
IHC:	User Optimized
IP:	User Optimized
WB:	1:1000


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	1X PBS, pH 7.4
Preservative:	0.09% (w/v) Sodium Azide
Stabilizer:	50% (v/v) Glycerol

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of mouse anti-Rhodopsin antibody. Lane 1:
 Human Cell lines. Primary antibody: Rhodopsin antibody at
 1:1000 for overnight at 4°C. Secondary antibody: Goat anti-
 mouse IgG HRP secondary antibody at 1:10,000 for 45 min at
 RT. Block: 5% Biotin overnight 4°C. Predicted/Observed size:
 38.8 kDa/40kD. Other band(s): none.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.