

Datasheet for 200-301-MX7

hFcRn (ADM31) Antibody

Overview

Description:	Anti-hFcRn (MOUSE) Monoclonal Antibody - 200-301-MX7
Item No.:	200-301-MX7
Size:	100 µg
Applications:	ELISA, FC, IF
Reactivity:	Human
Host Species:	Mouse

Product Details

Background:	The MHC class I-like Fc receptor (FcRn) functions as an intracellular trafficking receptor, uniquely accountable for prolonging the serum half-life of IgG subclass antibodies and facilitating their passage across cellular barriers. Through these actions, FcRn significantly impacts various aspects of antibody biology and pathobiology. Its crucial role in regulating IgG pharmacokinetics has been exploited in the development of therapeutic antibodies and related biological treatments. Additionally, FcRn is involved in the transport of serum albumin, contributing to the improved pharmacokinetic characteristics of albumin-conjugated therapies. However, a lack of dependable serological tools targeting human FcRn has limited the comprehension of its functions and potential therapeutic uses.
Synonyms:	FCRN antibody, alpha-chain antibody, FcRn antibody, Fc fragment of IgG receptor and transporter antibody, Fc receptor, IgG, alpha chain transporter antibody, FCGRT antibody, Fcgrt antibody, IgG receptor FcRn large subunit p51, Neonatal Fc receptor, FCGRT
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	ADM31
Format:	IgG2b

Target Details

Gene Name:	FCGRT
Reactivity:	Human

Immunogen:	Antibodies were generated by immunizing mouse FcRn-deficient mice with spleen cells from mice transgenic for human FcRn.
Purity/Specificity:	This protein G purified mouse monoclonal antibody was raised via hybridoma and recognizes hFcRn. Antibody isotype IgG2b lambda.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P55899• SDS

Application Details

Tested Applications:	ELISA, FC, IF
Application Note:	This monoclonal anti-hFcRn (ADM31) antibody is unconjugated and is suitable for the detection of FcRn (Fc receptor) in human samples. The FcRn antibody has been tested for Cell ELISA, ELISA, FACS and IF. For each application a titration should be performed to determine the optimal concentration.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:200 - 1:400
FC:	1.2 µg/10 ⁶ cells
IF:	User Optimized


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.068 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Stabilizer:	50% (v/v) Glycerol

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Immunofluorescence Microscopy


Immunofluorescence of Mouse Anti-hFcRn (ADM31) Antibody.

Cells: huSSECTM cells overexpressing hFcRn-EGFP.

Primary Antibody: Anti-hFcRn (ADM31) Antibody at 15µg/mL.

Secondary Antibody: Rabbit Anti-Mouse IgG Texas Red Conjugated.

A: DAPI-nuclear counterstain, B: hFcRn-GFP, C: Texas Red anti-FcRn (ADM31), D: merge.



Immunofluorescence Microscopy

Immunofluorescence of Mouse Anti-hFcRn (ADM31) Antibody.

Cells: huSSECTM cells.

Fixation: 4% PFA.

Permeabilization: PBS + 0.3% Triton X-100 + 1% goat serum.

Primary Antibody: Anti-hFcRn (ADM31) Antibody at 15µg/mL overnight at 4°C.

Secondary Antibody: Rabbit Anti-Mouse IgG Texas Red Conjugated (p/n 610-409-C46) 1µg/mL for 1hr at room temperature.

Location: Intracellular.


Flow Cytometry


Flow Cytometry of Mouse Anti-hFcRn (ADM31) Antibody.

Cells: huSSECTM cells.


Primary Antibody: Anti-hFcRn (ADM31) Antibody 1.2µg incubated on ice (4°C) for 60mins.

Secondary Antibody: Rabbit Anti-Mouse IgG Texas Red Conjugated (p/n 610-409-C46) 1.2µg incubated on ice (4°C) for 60mins protected from light.




ELISA

Identification of monoclonal Abs that block albumin binding to hFcRn. Representative sensorgrams showing binding of titrated amounts of monomeric hFcRn injected over immobilized ADM31 at pH7.4. Fig 3. PMID: 24764301.


ELISA

The monoclonal Abs bind in a species-dependent manner to FcRn. ELISA binding of human, macaque, pig, dog, mouse, and rat FcRn toward ADM31. Fig 5. PMID: 24764301.

References

- Sand KMK, Dalhus B, Christianson GJ, Bern M, Foss S, Cameron J, Sleep D, Bjoras M, et al. Dissection of the Neonatal Fc Receptor (FcRn)-Albumin Interface Using Mutagenesis and Anti-FcRn Albumin-blocking Antibodies. *J Biol Chem.* (2014)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.