

## Datasheet for 200-301-MX8C

# hFcRn (DVN24) Antibody

### Overview

<b>Description:</b>	Anti-hFcRn (MOUSE) Monoclonal Antibody - 200-301-MX8C
<b>Item No.:</b>	200-301-MX8C
<b>Size:</b>	100 µg
<b>Applications:</b>	ELISA, FC, IF
<b>Reactivity:</b>	Human
<b>Host Species:</b>	Mouse

### Product Details

<b>Background:</b>	The MHC class I-like Fc receptor (FcRn) is an intracellular trafficking Fc receptor that is uniquely responsible for the extended serum half-life of antibodies of the IgG subclass and their ability to transport across cellular barriers. By performing these functions, FcRn affects numerous facets of antibody biology and pathobiology. Its critical role in controlling IgG pharmacokinetics has been leveraged for the design of therapeutic antibodies and related biologics. FcRn also traffics serum albumin and is responsible for the enhanced pharmacokinetic properties of albumin-conjugated therapeutics.
<b>Synonyms:</b>	FCRN antibody, alpha-chain antibody, FcRn antibody, Fc fragment of IgG receptor and transporter antibody, Fc receptor, IgG, alpha chain transporter antibody, FCGRT antibody, Fcgrt antibody, IgG receptor FcRn large subunit p51, Neonatal Fc receptor, FCGRT
<b>Host Species:</b>	Mouse
<b>Clonality:</b>	Monoclonal
<b>Clone ID:</b>	DVN24
<b>Format:</b>	IgG2a

### Target Details

<b>Gene Name:</b>	FCGRT
<b>Reactivity:</b>	Human
<b>Immunogen:</b>	Antibodies were generated by immunizing mouse FcRn-deficient mice with spleen cells from mice transgenic for human FcRn.

**Purity/Specificity:** This protein G purified mouse monoclonal antibody was raised via hybridoma and recognizes hFcRn. Antibody isotype IgG2a kappa.

**Relevant Links:**

- [UniProtKB - P55899](#)

## Application Details

**Tested Applications:** ELISA, FC, IF

**Application Note:** This monoclonal anti-hFcRn (DVN24) antibody is unconjugated and is suitable for the detection of FcRn (Fc receptor) in human samples. The FcRn antibody has been tested for Cell ELISA, ELISA, FACS and IF. For each application a titration should be performed to determine the optimal concentration.

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

**ELISA:** 1:200 - 1:400

**FC:** 1.2 µg/10<sup>6</sup> cells

**IF:** User Optimized

## Formulation

**Physical State:** Liquid (sterile filtered)

**Concentration:** 1.011 mg/mL by UV absorbance at 280 nm

**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Preservative:** None

**Stabilizer:** None


## Shipping & Handling

**Shipping Condition:** Wet Ice

**Storage Condition:** Store vial at 4° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing.

**Expiration:** Expiration date is one (1) year from date of receipt.

## Images




### Immunofluorescence Microscopy

Immunofluorescence of Mouse Anti-hFcRn (DVN24) Antibody.

Cells: huSSECTM cells over expressing hFcRn-EGFP.  
 Primary Antibody: Anti-hFcRn (DVN24) Antibody at 15µg/mL.

Secondary Antibody: Rabbit Anti-Mouse IgG Texas Red Conjugated.

A: DAPI-nuclear counterstain, B: hFcRn-GFP, C: Texas Red anti-FcRn (DVN24), D: merge.



### Immunofluorescence Microscopy

Immunofluorescence of Mouse Anti-hFcRn (DVN24) Antibody.

Cells: huSSECTM cells.

Fixation: 4% PFA.

Permeabilization: PBS + 0.3% Triton X-100 + 1% goat serum.  
 Primary Antibody: Anti-hFcRn (DVN24) Antibody at 15µg/mL overnight at 4°C.

Secondary Antibody: Rabbit Anti-Mouse IgG Texas Red Conjugated (p/n 610-409-C46) 1µg/mL for 1hr at room temperature.

Location: Intracellular.


### Flow Cytometry


Flow Cytometry of Mouse Anti-hFcRn (DVN24) Antibody.

Cells: huSSECTM cells.


Primary Antibody: Anti-hFcRn (DVN24) Antibody 1.2µg incubated on ice (4°C) for 60mins.

Secondary Antibody: Rabbit Anti-Mouse IgG Texas Red Conjugated (p/n 610-409-C46) 1.2µg incubated on ice (4°C) for 60mins protected from light.




**ELISA**

Identification of monoclonal Abs that block albumin binding to hFcRn. Representative sensorgrams showing binding of titrated amounts of monomeric hFcRn injected over immobilized DVN24 at pH7.4. Fig 3. PMID: 24764301.


**ELISA**

The monoclonal Abs bind in a species-dependent manner to FcRn. ELISA binding of human, macaque, pig, dog, mouse, and rat FcRn toward DVN24. Fig 5. PMID: 24764301.

## References

- Sand KMK, Dalhus B, Christianson GJ, Bern M, Foss S, Cameron J, Sleep D, Bjoras M, et al. Dissection of the Neonatal Fc Receptor (FcRn)-Albumin Interface Using Mutagenesis and Anti-FcRn Albumin-blocking Antibodies. *J Biol Chem.* (2014)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.