

Datasheet for 200-302-N62**CD8a Fluorescein Antibody****Overview**

Description:	Anti-CD8a (MOUSE) Fluorescein Conjugated Monoclonal Antibody - 200-302-N62
Item No.:	200-302-N62
Size:	500 µL
Applications:	FC
Reactivity:	Human
Host Species:	Mouse

Product Details

Background:	The OKT8 antibody is specific for the 32-34 kDa alpha chain of human CD8, known as CD8a or CD8 alpha. CD8a can form a homodimer (CD8 alpha-alpha), but is more commonly expressed as a heterodimer with a second chain known as CD8b or CD8 beta. CD8 acts as a co-receptor for antigen recognition and subsequent T cell activation that is initiated upon binding of the T cell receptor (TCR) to antigen-bearing MHC Class I molecules. The cytoplasmic domains of CD8 provide binding sites for the tyrosine kinase lck, facilitating intracellular signaling events that lead to T cell activation, development, and cytotoxic effector functions. CD8+ cytotoxic T cells (CTLs) play an important role in inducing cell death of tumor cells, as well as cells infected by virus, bacteria or parasites. The OKT8 antibody is widely used as a phenotypic marker for CD8 on cytotoxic T cells, thymocytes, as well as on certain cell types that do not also express the TCR, including some NK cells and lymphoid dendritic cells.
Synonyms:	T-cell surface glycoprotein CD8 alpha chain, T-lymphocyte differentiation antigen T8/Leu-2, CD8a
Host Species:	Mouse
Conjugate:	Fluorescein (FITC)
Clonality:	Monoclonal
Clone ID:	OKT8
Format:	IgG2a
F/P Ratio:	2-8

Target Details

Gene Name:	CD8A
Reactivity:	Human
Immunogen:	Anti-CD8a Antibody (Monoclonal) was produced by repeated immunizations with CD8a antigen.
Purity/Specificity:	Fluorescein conjugated CD8a Monoclonal Antibody was purified from tissue culture supernatant via affinity chromatography and is directed against human CD8a. Reactivity is observed against human CD8a. Cross reactivity with CD8a from other sources has not been tested. Anti-CD8a is conjugated with FITC under optimal conditions and the solution is free of unconjugated FITC.
Relevant Links:	• UniProtKB - P01732 • NCBI - NP_001139345.1 • GeneID - 925

Application Details

Tested Applications:	FC
Application Note:	Anti-CD8a is tested for Flow Cytometry and is useful for Immunohistochemistry. Researchers should determine optimal titers for applications that are not stated.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
FC:	5µL/test/1x10e5 to 1x10e8 cells
IHC:	User Optimized
IP:	User Optimized


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.025mg/mL Sufficient to run approximately 100 tests
Buffer:	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.09% (w/v) Sodium Azide
Stabilizer:	0.1% Gelatin

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. DO NOT FREEZE. This product is light sensitive.
Expiration:	Expiration date is six (6) months from date of receipt.

Images



Flow Cytometry

Flow Cytometry of Mouse anti-CD8a Fluorescein Conjugated Monoclonal Antibody. Cells: human peripheral blood lymphocytes. Stimulation: none. Antibody: (Dotted line) FITC Mouse IgG2a isotype control; (solid line) Fluorescein Anti-CD8a mouse antibody using 5 ul (0.125µg).

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.