

Datasheet for 200-326-N25**CD25 APC Antibody****Overview**

Description:	Anti-CD25 (MOUSE) Allophycocyanin Conjugated Monoclonal Antibody - 200-326-N25
Item No.:	200-326-N25
Size:	500 µL
Applications:	FC
Reactivity:	Human
Host Species:	Mouse

Product Details

Background:	The BC96 antibody is specific for human CD25, a 55 kDa surface protein also known as the Interleukin-2 Receptor alpha chain, or IL-2R alpha. CD25 may bind IL-2 by itself, although with low affinity and without induction of cell signaling. CD25 is also expressed within a high-affinity complex, along with the IL-2R beta chain (CD122) and the common gamma chain (CD132), to form a signaling receptor complex. Signaling via CD25 may modulate Th17 and Treg cell differentiation, and has been shown to play a role in the function of dendritic cells. The BC96 antibody may be used as a marker for expression of CD25 during T and B cell development, on activated mature T and B cells, and for its expression at high levels on natural T regulatory cells (nTreg cells) in the thymus or periphery, and on induced peripheral Tregs.
Synonyms:	Interleukin-2 receptor subunit alpha, IL-2 receptor subunit alpha, IL-2-RA, IL-2R subunit alpha, IL2-RA, TAC antigen, p55
Host Species:	Mouse
Conjugate:	Allophycocyanine (APC)
Clonality:	Monoclonal
Clone ID:	BC96
Format:	IgG1
F/P Ratio:	1-2

Target Details

Gene Name:	IL2RA
-------------------	-------

Reactivity:	Human
Immunogen:	Anti-CD25 Antibody (Monoclonal) was produced by repeated immunizations with CD25 antigen.
Purity/Specificity:	Allophycocyanin conjugated CD25 Monoclonal Antibody was purified from tissue culture supernatant via affinity chromatography and is directed against human CD25. Reactivity is observed against human CD25, Baboon, Chimpanzee, Cynomolgus, and Rhesus. Cross reactivity with CD25 from other sources has not been tested. Anti-CD25 is conjugated with APC under optimal conditions and the solution is free of unconjugated APC.
Relevant Links:	• UniProtKB - P01589 • NCBI - NP_000408.1 • GeneID - 3559

Application Details

Tested Applications:	FC
Application Note:	Anti-CD25 is tested for Flow Cytometry. Researchers should determine optimal titers for applications that are not stated.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
FC:	5µL/test/1x10e5 to 1x10e8 cells


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	0.025 mg/mL Sufficient to run approximately 100 tests
Buffer:	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.09% (w/v) Sodium Azide
Stabilizer:	0.1% Gelatin

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. Dilute only prior to immediate use. This product is stable at 4° C as an undiluted liquid. Use subdued lighting during handling and incubation of cells prior to analysis. Store reagent in the dark. DO NOT FREEZE.
Expiration:	Expiration date is six (6) months from date of receipt.

Images



Flow Cytometry

Flow Cytometry of Mouse anti-CD25 Allophycocyanin Conjugated Monoclonal Antibody. Cells: human peripheral blood lymphocytes. Stimulation: none. Antibody: (LEFT) APC Mouse IgG1 kappa isotype control and FITC Anti-CD4 using 5 μ L; (RIGHT) Allophycocyanin Anti-CD25 mouse antibody and FITC Anti-CD4 using 5 μ L.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.