

Datasheet for 200-401-033S

Ovalbumin Antibody

Overview

| | |
|----------------------|---|
| Description: | Anti-Ovalbumin (Hen Egg White) (RABBIT) Antibody - 200-401-033S |
| Item No.: | 200-401-033S |
| Size: | 25 µL |
| Applications: | WB, Biochemical Assay, ELISA, IHC |
| Reactivity: | Chicken |
| Host Species: | Rabbit |

Product Details

Background: Anti-Ovalbumin Antibody recognizes ovalbumin that is the main protein found in egg white. Ovalbumin makes up 55-65% of the total protein. Ovalbumin displays sequence and three-dimensional homology to the serpin superfamily, but unlike most serpins, it does not undergo a conformational change upon proteolytic cleavage and is not a serine protease inhibitor. The function of ovalbumin is still unknown, although it is presumed to be a storage protein for tissue-specific, steroid hormone-induced gene expression. Ovalbumin is the second most common food allergen in infants and is widely used as an antigen to induce IgE-mediated food allergy animal models, allergic asthma models, and atopic dermatitis models. OVA was used as a model antigen to study the efficacy of cancer vaccines using nanoparticles, where OVA (instead of a tumor antigen) provides a convenient method to evaluate B-cell response. Ovalbumin is a key protein for biochemical investigations, processes, allergen disease models, and vaccine studies. It can be used as a control, blocking reagent, or antibody in assays such as ELISA, SDS-PAGE, IHC, Western Blot, and size exclusion chromatography.

| | |
|----------------------|---|
| Synonyms: | rabbit anti-Ovalbumin Antibody, Egg albumin antibody, Plakalbumin, Allergen Gal d II, Allergen=Gal d 2, Hen Egg White |
| Host Species: | Rabbit |
| Clonality: | Polyclonal |
| Format: | IgG |

Target Details

| | |
|-------------------|-----------|
| Gene Name: | SERPINB14 |
|-------------------|-----------|

| | |
|----------------------------|---|
| Reactivity: | Chicken |
| Immunogen Type: | Native Protein |
| Immunogen: | Ovalbumin [Hen Egg White] |
| Purity/Specificity: | Rabbit Anti-Ovalbumin is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as purified and partially purified Ovalbumin [Hen Egg White]. Cross reactivity against Ovalbumin from other sources is unknown. |
| Relevant Links: | <ul style="list-style-type: none">• UniProtKB - P01012• NCBI - P01012.2• GeneID - 396058 |

Application Details

| | |
|--------------------------------|--|
| Tested Applications: | WB |
| Suggested Applications: | Biochemical Assay, ELISA, IHC (Based on references) |
| Application Note: | Anti-Ovalbumin Antibody has been tested in Western Blot. Although not tested this product could be useful in assays such as ELISA, IHC, and IP. Specific conditions for reactivity and signal detection should be optimized by the end user. |
| Assay Dilutions: | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| ELISA: | 1:25,000 |
| IHC: | 1:300 - 1,2000 |
| IP: | 1:100 |
| WB: | 1:500 - 1:2,000 |

Formulation

| | |
|------------------------|--|
| Physical State: | Liquid (sterile filtered) |
| Concentration: | 1.0 mg/ml by UV absorbance at 280 nm |
| Buffer: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Preservative: | 0.01% (w/v) Sodium Azide |
| Stabilizer: | None |

Shipping & Handling

Shipping Condition: Dry Ice


Storage Condition: Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.

Expiration: Expiration date is three (3) months from date of receipt.


Images



Control



Canna




OVA



OVA-Canna

Immunohistochemistry

Immunohistochemical Analysis Using Anti-Ovalbumin Antibody. After the evaluation of anaphylactic symptoms on day 28, the liver was collected, fixed, and used to prepare frozen sections. The samples were incubated overnight with a primary rabbit polyclonal ovalbumin antibody (p/n 200-401-033). The images are representative of each group. Immunohistochemical staining of liver sections was used to determine the effects of canna starch on OVA localization (400×). Fig 2. PMID: 38397452.



Western Blot


Cross-presentation of OVA antigen carried by the subtypes of particles.

(A) Scheme of the construct containing the myristoylation and palmitoylation sequences fused to OVA, myc tag, P2A cleavage site and puromycin resistance gene for selection, which was used to create the EO771 myr/palm-OVA stable cells.

(B) Analysis of EVs/ENPs from 20×10^6 EO771-m/p-OVA by western Blot. Four known amounts of recombinant OVA were loaded on the same gel to allow quantification of OVA. Blots were revealed with a polyclonal antibody against OVA. One representative blot out of 6. Arrowheads indicate the positions of myr/palm-OVA (upper band) and recombinant OVA (lower band).


(C) Quantification of OVA in the different EVs/ENPs, was done on the western blot images as compared to the OVA dose-response curve. Graphs show ng OVA/10⁶ cells (left) or ng OVA/particles (right). Friedman test with Dunn's multiple comparison was performed. n = 6.

Figure 6. PMID: 38073509




Figure

Sensograms of two spots in response to an Anti-IgG injection and two regeneration steps in sequence. The two sensograms correspond to the reference spot with anti-ovalbumin antibody (R, black curve) and the signal spot with human IgG antibody (M, blue curve). Then 10 min regeneration steps are performed to recover the starting conditions (before Anti-IgG injection, I marker). W indicates the washing steps. Fig 4. PMID: 30044392



Western Blot

Western Blot of Rabbit anti-OVALBUMIN (Hen Egg White) antibody. Lane 1: Reduced Ovalbumin. Load: 50 ng. Primary antibody: Ovalbumin antibody at 1:1,000 overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody (p/n 611-103-122) at 1:40,000 for 30 min at RT. Block: Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) for 30 min at RT. Predicted/Observed size: 45 kDa, 45 kDa for Ovalbumin.



Western Blot

Western Blot of Rabbit anti-Ovalbumin antibody. Lane 1: Ovalbumin protein - Reduced [$\sim 1 \mu\text{g}$]. Lane 2: Ovalbumin protein - Reduced [$0.25 \mu\text{g}$]. Lane 3: Ovalbumin protein - Non-reduced [$\sim 1 \mu\text{g}$]. Lane 4: Ovalbumin protein - Non-reduced [$0.25 \mu\text{g}$]. Primary antibody: Ovalbumin antibody at 1:5000 for overnight at 4°C. Secondary antibody: Atto 425 conjugated goat anti-rabbit secondary antibody at 1:10,000 for 1.5 hr at RT. Block: MB-070 buffer overnight at 4°C. Predicted/Observed size: 42.9 kDa, ~ 36 kDa for ovalbumin. Other band(s): none.

References

- Koida A et al. Canna Starch Improves Intestinal Barrier Function, Inhibits Allergen Uptake, and Suppresses Anaphylactic Symptoms in Ovalbumin-Induced Food Allergy in Mice. *Biomolecules*. (2024)
- Cocozza F et al. Extracellular vesicles and co-isolated endogenous retroviruses from murine cancer cells differentially affect dendritic cells. *EMBO J*. (2023)
- Sinibaldi A et al. Label-free monitoring of human IgG/anti-IgG recognition using bloch surface waves on 1D photonic crystals. *Biosensors (Basel)*. (2018)
- Dufresne-Martin G et al. Peptide mass fingerprinting by matrix-assisted laser desorption ionization mass spectrometry of proteins detected by immunostaining on nitrocellulose. *Proteomics*. (2005)
- Kathryn L Brogan et al. Influence of antibody immobilization strategy on molecular recognition force microscopy measurements. *Langmuir*. (2005)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.