

Datasheet for 200-401-698S**Jagged 1 Antibody****Overview**

| | |
|----------------------|--|
| Description: | Anti-Jagged-1 (RABBIT) Antibody - 200-401-698S |
| Item No.: | 200-401-698S |
| Size: | 25 µL |
| Applications: | ELISA, IF, IHC, WB |
| Reactivity: | Human, Mouse |
| Host Species: | Rabbit |

Product Details

| | |
|----------------------|--|
| Background: | Anti-Jagged 1 Antibody recognizes the Jagged 1 protein encoded by the JAG1 gene, that is the human homolog of the Drosophila jagged protein. Human Jagged 1 is the ligand for multiple Notch receptors and mediation of Notch signaling. Jagged 1 signaling through Notch 1 has also been shown to play a role in hematopoiesis. Mutations that alter the Jagged 1 protein cause Alagille syndrome, deafness, congenital heart defects, and posterior embryotoxon. Anti-Jagged-1 Antibody is useful for researchers interested in Notch activation, Growth Factor activities, and Cardiovascular health. |
| Synonyms: | rabbit anti-Jagged 1 Antibody, rabbit anti-Jagged1 Antibody, rabbit anti-Jagged-1 Antibody, Ser 1 antibody, AGS antibody, AHD antibody, AWS antibody, CD 339 antibody, CD339 antibody, CD339 antigen antibody, Headturner antibody, HJ1 antibody, Htu antibody |
| Host Species: | Rabbit |
| Clonality: | Polyclonal |
| Format: | IgG |

Target Details

| | |
|------------------------|--------------------|
| Gene Name: | JAG1 |
| Reactivity: | Human, Mouse |
| Immunogen Type: | Conjugated Peptide |

| | |
|----------------------------|--|
| Immunogen: | This protein A purified Jagged 1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to amino acids near the N-terminus of human Jagged-1 protein. |
| Purity/Specificity: | This protein A purified antibody is directed against human Jagged-1 protein. The product was purified from mono-specific antiserum by affinity chromatography. A BLAST analysis was used to suggest cross reactivity with Jagged-1 protein from human, chimpanzee, rat and mouse based on 100% homology with the immunizing sequence. Partial reactivity is expected against dog (81%) and <i>Xenopus laevis</i> (85%) based on partial sequence homologies as indicated. Reactivity against homologues from other sources is not known. |
| Relevant Links: | <ul style="list-style-type: none">• UniProtKB - P78504• GeneID - 182• NCBI - 4557679 |

Application Details

| | |
|-----------------------------|---|
| Tested Applications: | ELISA, IF, IHC, WB |
| Application Note: | This protein A purified antibody has been tested for use in ELISA, immunohistochemistry, immunofluorescence microscopy and western blot. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 150 kDa in size corresponding to Jagged-1 in mouse liver, human liver and human lung whole cell lysates. |
| Assay Dilutions: | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| ELISA: | 1:20,000 - 1:100,000 |
| IF: | 1:200 - 1:1,000 |
| IHC: | 1:100 - 1:500 |
| WB: | 1:500 - 1:2,000 |


Formulation

| | |
|------------------------|--|
| Physical State: | Liquid (sterile filtered) |
| Concentration: | 1.1 mg/mL by UV absorbance at 280 nm |
| Buffer: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Preservative: | 0.01% (w/v) Sodium Azide |
| Stabilizer: | None |

Shipping & Handling


| | |
|----------------------------|--|
| Shipping Condition: | Dry Ice |
| Storage Condition: | Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing. |
| Expiration: | Expiration date is one (1) year from date of receipt. |

Images



Immunocytochemistry

(a) Representative confocal microscopy shows CD44, CD24 and cleaved notch (NICD) in a population of drug naive MDA-MB-231. Yellow arrows indicate CD44^{hi}CD24^{lo} (M) population of cells and the white arrows indicate the CD44^{hi}CD24^{hi} (E/M) cells. Histogram (right panel) shows quantification of NICD in the distinct phenotype populations (M versus E/M). N = 3 biological replicates. (b) Schematic describes the experimental protocol to generate drug-tolerant cells (DTCs) parental MDA-MB-231 cells were treated with docetaxel at 100 nM (20× the IC₅₀) and subsequently selected by substrate re-attachment and acute population outgrowth. (c) Representative confocal microscopy shows CD44, CD24 and NICD in the MDA-MB-231 parent and DTC populations. Right panel shows quantification of fluorescence intensity of each signal determined by at least 25 individual fields. N = 3 biological replicates. (d) Representative confocal microscopy shows Jagged and Delta expression in MDA-MB-231 parent and DTC. DAPI nuclear stain (blue). N = 3 biological replicates. Figure provided by CiteAb. Source: J R Soc Interface, PMID: 27170649.



Western Blot


Western blot Anti-Jagged-1 (RABBIT) Antibody. Human pluripotent stem cells (lanes 1 and 2), definite endoderm cells (lanes 3 and 4), pancreatic endoderm cells (lanes 5 and 6), and pancreatic progenitors (lanes 7 and 8). Primary Antibody: Anti-Jagged-1 used at a 1:1000 (30sec exposure) overnight at 4°C. Beta-actin served as loading control at 1:2000 at RT for 1hr.

Secondary Antibody: donkey anti-rabbit HRP conjugated antibody or donkey anti-mouse HRP conjugated antibody diluted 1:5000 for 1h at RT.

Independently Validated by antibodies-online GmbH (p/n ABIN1043767/ ABIN129524) courtesy of Ulm University Hospital.


Immunohistochemistry


Immunohistochemical staining of human cervical cancer tissue (40X magnification) using Rockland's Protein A purified anti-Jagged-1 antibody. Tissue was fixed with formalin and embedded in paraffin. Hematoxylin was used to counter-stain cells. A 1:100 dilution of primary antibody was used. Personal Communication. Martin Kast Laboratory.



Immunofluorescence Microscopy

Immunofluorescence microscopy using Rockland's Protein A purified anti-Jagged-1 antibody of human corneal epithelial cells. Primary antibody was used at a 1:500 dilution. The Jagged1 (green staining) is localized to the cytoplasm and is consistent with reports in the literature. The nucleus is stained with Bis benzamine (blue). Personal Communication. Aihua Ma, University of Cardiff.



**Western Blot**

Western Blot of Rabbit Anti-Jagged 1 Antibody. Lane 1: Opal Prestained MW marker (p/n MB-210-0500). Lane 2: Mouse Liver Whole Cell Lysate [10 μ g] (p/n W10-000-T020). Lane 3: Human Liver Whole Cell Lysate [10 μ g]. Lane 4: Human Lung Whole Cell Lysate [10 μ g]. Primary Antibody: Anti-Jagged 1 at 1:1000 overnight at 2-8°C. Secondary Antibody: Goat Anti-Rabbit IgG Peroxidase Conjugated (p/n 611-103-122) at 1:70000 for 30mins at RT. Blocking Buffer: BlockOut Buffer (p/n MB-073) for 1hr RT. Predicted Molecular Weight: 113kDa. Exposure: 30 sec.

References

- Boareto et al. Notch-Jagged signalling can give rise to clusters of cells exhibiting a hybrid epithelial/mesenchymal phenotype. *Journal of the Royal Society Interface* (2016)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.