Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 200-401-A65

TLR10 Antibody

Overview

Description:	Anti-TLR10 (RABBIT) Antibody - 200-401-A65
Item No.:	200-401-A65
Size:	100 µg
Applications:	ELISA, IF, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Toll-like receptors (TLRs) are signaling molecules that recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. These proteins act through adaptor molecules such as MyD88 and TIRAP to activate various kinases and transcription factors. TLR10 is highly homologous to TLRs 1 and 6 and is most highly expressed in lymphoid tissues. Most recently genetic variation in TLR10 has been associated with contributing to asthma risk. It should be noted that TLR10 exists in mice only as a pseudogene.
Synonyms:	CD290 antibody, CD290 antigen antibody, Toll like receptor 10 antibody
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	TLR10
Reactivity:	Human
Immunogen Type:	Conjugated Peptide
Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region of human toll-like receptor 10 protein.

Purity/Specificity:	This antibody is directed against human TLR10 protein. This product is affinity chromatography purified via peptide column. This antibody is reactive with TLR10 from human sources. Cross reactivity with TLR10 from other sources has not been determined.
Relevant Links:	• NCBI - AAY78491.1 • UniProtKB - Q9BXR5 • GeneID - 81793

Application Details

Tested Applications:	ELISA, IF, WB
Application Note:	This IgG fraction antibody has been tested for use in ELISA, immunofluorescence, and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band at ~75 kDa in size corresponding to TLR10 by western blotting in the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:40,000
IF:	2 µg/mL
WB:	1:500 - 1:3,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot
Western Blot of Rabbit Anti-TLR10 Antibody. Lane 1: human lymph node whole cell lysate using 0.5 µg/ml. Lane 2: human lymph node whole cell lysate using 1.0 µg/ml. Load: 35 µg per lane. Primary antibody: TLR-10 antibody overnight at 4°C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 95 kDa, ~75 kDa for TLR10. Other band(s): none.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.