

Datasheet for 200-401-A68**TLR1 Antibody****Overview**

Description:	Anti-TLR1 (RABBIT) Antibody - 200-401-A68
Item No.:	200-401-A68
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Toll-like receptors (TLRs) are evolutionarily conserved pattern-recognition molecules resembling the toll proteins that mediate antimicrobial responses in <i>Drosophila</i> . These proteins recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. The TLRs act through adaptor molecules such as MyD88 and TIRAP to activate various kinases and transcription factors so the organism can respond to potential infection. TLR1 is co-expressed with TLR2 on myeloid cells of the innate immune systems in lymphoid tissue such as monocytes and dendritic cells where they form heterodimers that can recognize triacylated lipoproteins.
Synonyms:	CD281 antigen antibody, rsc786 antibody, Similar to <i>Drosophila</i> Toll protein antibody, TIL antibody, Toll (<i>Drosophila</i>) homolog antibody, Toll like receptor 1 Precursor antibody
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	TLR1
Reactivity:	Human
Immunogen Type:	Conjugated Peptide

Immunogen:	This affinity purified antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to a region near the amino terminus of human toll-like receptor 1 (TLR1) protein.
Purity/Specificity:	This IgG fraction antibody is directed against human TLR1 protein. This product is affinity chromatography purified via peptide column. This antibody is reactive with TLR1 from human and mouse sources. Cross reactivity with TLR1 from other sources has not been determined.
Relevant Links:	• NCBI - CAG38593.1 • UniProtKB - Q15399 • GeneID - 7096

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	This IgG fraction antibody has been tested for use in ELISA, immunohistochemistry, immunofluorescence, and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band at ~90 kDa in size corresponding to TLR1 by western blotting in the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000 - 1:40,000
IF:	20 µg/mL
IHC:	10 µg/ml
WB:	1 - 4 µg/ml

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Rabbit Anti-TLR1 Antibody. Lane 1: mouse spleen tissue whole cell lysate using 1.0 ug/ml. Lane 2: mouse spleen tissue whole cell lysate using 2.0 ug/ml. Lane 3: mouse spleen tissue whole cell lysate using 4.0 ug/ml. Load: 10 µg per lane. Primary antibody: TLR-1 antibody overnight at 4°C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 90 kDa, ~90 kDa for TLR 1. Other band(s): none.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.