

Datasheet for 200-401-B48**CENP-Q Antibody****Overview**

Description:	Anti-CENP-Q (RABBIT) Antibody - 200-401-B48
Item No.:	200-401-B48
Size:	100 µg
Applications:	ELISA, IF, Multiplex, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	This antibody is designed, produced, and validated as part of a collaboration between Rockland and the National Cancer Institute (NCI) and is suitable for Cancer, Immunology and Nuclear Signaling research. Cenp-Q (also known as centromere protein Q or CENPQ) is a nuclear/centromeric protein that is one of the critical components that constitutes the CENP-O complex at the kinetochores and appears to stabilize PBIP1/CENP-U(50)/MLF1IP in the complex. This complex is important for proper recruitment of polo-like kinase 1 (Plk1) to the mitotic kinetochores. A failure in this process results in improper microtubule attachment to the kinetochores and chromosome missegregation that ultimately lead to aneuploidy.
Synonyms:	rabbit anti-CENP-Q Antibody, CenpQ, centromere protein Q, CENP Q
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	CENPQ
Reactivity:	Human
Immunogen Type:	Recombinant Protein
Immunogen:	This protein A purified antibody was prepared from whole rabbit serum produced by repeated immunizations with full-length human CENP-Q recombinant protein.

Purity/Specificity: This product was protein A purified from monospecific antiserum by immunoaffinity chromatography using protein A coupled to agarose beads. This antibody is specific for human CENP-Q protein. A BLAST analysis of the full length sequence was used to suggest partial cross-reactivity with CENP-Q based on the following percentage homologies: macaque (91%), horse (76%), bovine (74%), dog (73%), swine (71%), mouse (65%) and rat (60%). Cross-reactivity with CENP-Q from other sources has not been determined.

Relevant Links:

- [NCBI - 40068061](#)
- [UniProtKB - Q7L2Z9](#)
- [GeneID - 55166](#)

Application Details

Tested Applications: ELISA, IF, Multiplex, WB

Application Note: This protein A purified antibody has been tested for use in ELISA, immunofluorescence microscopy and western blotting. Specific conditions for reactivity should be optimized by the end user. Expect a band approximately 26-31 kDa in size corresponding to human CENP-Q by western blotting in the appropriate cell lysate or extract.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:5,000 1:20,000

IF: User Optimized

WB: 1:100 - 1:500

Formulation

Physical State: Liquid (sterile filtered)

Concentration: 1.15 mg/mL by UV absorbance at 280 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: 0.01% (w/v) Sodium Azide

Stabilizer: None


Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.


Images



Western Blot


The Plk1 interacts with and phosphorylates CENP-Q via the PBIP1 T78 motif. A and B, HeLa cells transfected with the indicated constructs were subjected to immunoprecipitation analyses. Where indicated, cells were treated with nocodazole for 18 h and BI 2536 for 4 h before harvest. Immunoprecipitates were reacted with λ phosphatase to convert all the phosphorylated, slow-migrating forms to non-phosphorylated form for easy detection by immunoblotting analysis. Bracket, hyperphosphorylated PBIP1 forms. C, HeLa cells expressing the indicated RNAi-insensitive constructs were depleted of control luciferase (si-Luc) or endogenous PBIP1 (si-PBIP1), and then treated with nocodazole for 18 h before harvest. Samples were immunoprecipitated with control IgG (first lane) or anti-CENP-Q antibody and the immunoprecipitates were treated with λ phosphatase before immunoblotting analyses. Asterisk, a cross-reacting protein. Note that cells depleted of PBIP1 exhibited a greatly diminished level of CENP-Q immunoprecipitates due to its instability in the absence of PBIP1 (lane 2). D and E, The cells used in C were depleted of endogenous PBIP1 (si-PBIP1), and then either left untreated or treated with nocodazole before harvest. Samples were separated 10% SDS-PAGE and immunoblotted. Transfected CENP-Q (no tag) in lane 1 denotes the position of endogenous CENP-Q. Note that the T78-dependent PBIP1 function is required for mitotic-specific phosphorylation of CENP-Q (D), but not for other Plk1 substrates, such as Bub1, BubR1, and Cdc25C (E). Asterisks, cross-reacting proteins. Fig 4.

PMID: 21454580



Immunofluorescence Microscopy

Immunofluorescence microscopy using Rockland's protein A purified anti-CENP-Q antibody shows detection of endogenous CENP-Q in HeLa whole cell lysate. Primary antibody was used at 1:100 followed by secondary antibody diluted 1:150. Red punctate anti-CENP-Q signal colocalizes in overlay images with green punctate anti-CREST signals at the kinetochores (attached points of sister chromatids). Visible are colocalized CENP-Q and CREST signal at various stages of the cell cycle as indicated from interphase to the end of mitosis. Nuclei are counter stained with bisbenzamide. Personal Communication, Kyung S. Lee, CCR-NCI, Bethesda, MD



Western Blot

Western blot using Rockland's protein A purified anti-CENP-Q antibody shows detection of endogenous CENP-Q in a HeLa whole cell lysate (lane 1, arrowhead). The blot was incubated for 1.5 hours at room temperature using the primary antibody diluted to 0.5 µg/mL, followed by washes and incubation with to the secondary antibody. Lane 1: Lysates from HeLa cells transfected with control sh-virus wherein the expression of CENP-Q is expected to not to alter. Lane 2: Lysates from HeLa cells transfected with Cenp-Q sh-virus wherein the expression of CENP-Q is knocked down significantly to a level where it is not being detected at all under the tested condition/WB exposure time. Personal Communication, Kyung S. Lee, CCR-NCI, Bethesda, MD.

References

- Kang YH, Park CH, Kim TS, Soung NK, Bang JK, Kim BY, Park JE, Lee KS. Mammalian polo-like kinase 1-dependent regulation of the PBIP1-CENP-Q complex at kinetochores. *Journal of Biological Chemistry* (2011)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.