

**Datasheet for 200-401-BF2****Flip Antibody****Overview**

|                      |   |
|----------------------|---|
| <b>Description:</b>  | Anti-FLIP (RABBIT) Antibody - 200-401-BF2 |
| <b>Item No.:</b>     | 200-401-BF2                               |
| <b>Size:</b>         | 100 µg                                    |
| <b>Applications:</b> | ELISA, IHC, WB                            |
| <b>Reactivity:</b>   | Mouse                                     |
| <b>Host Species:</b> | Rabbit                                    |

**Product Details**

|                      |  |
|----------------------|--|
| <b>Background:</b>   | Apoptosis is related to many diseases and induced by a family of cell death receptors and their ligands. Cell death signals are transduced by death domain (DD) containing adapter molecules and members of the ICE/CED-3 protease family. Caspases-8 (FLICE) and -10 (FLICE2) are two pivotal members in the ICE/CED-3 protease family. FLICE-inhibitory proteins were identified in virus and human and designated v-FLIPs and FLIP respectively. The human FLIP was also cloned by several labs independently and termed Casper, I-FLICE, FLAME-1, CASH, CLARP and MRIT. FLIP contains two death effector domains (DEDs) and a caspase-like domain. FLIP interacts with adapter protein FADD and caspase-8 and 10, and potently inhibits apoptosis induced by all known death receptors CD95, DR3, TRAIL-R and TNFR1. |
| <b>Synonyms:</b>     | FLIP Antibody, CASH, FLIP, MRIT, CLARP, FLAME, Casper, FLAME1, c-FLIP, FLAME-1, I-FLICE, c-FLIPL, c-FLIPR, c-FLIPS, CASP8AP1, CASP8 and FADD-like apoptosis regulator, Caspase homolog, Caspase-eight-related protein, Caspase-like apoptosis regulatory protein, Cellular FLICE-like inhibitory protein, FADD-like antiapoptotic molecule 1, Inhibitor of FLICE, MACH-related inducer of toxicity, Usurpin, Cflar   |
| <b>Host Species:</b> | Rabbit   |
| <b>Clonality:</b>    | Polyclonal   |
| <b>Format:</b>       | IgG  |

**Target Details**

|                    |       |
|--------------------|-------|
| <b>Gene Name:</b>  | Cflar |
| <b>Reactivity:</b> | Mouse |

|                            |   |
|----------------------------|---|
| <b>Immunogen Type:</b>     | Conjugated Peptide  |
| <b>Immunogen:</b>          | Anti-FLIP antibody was prepared from whole rabbit serum produced by repeated immunizations with a 17 amino acid peptide near the C-terminus of mouse FLIP.                    |
| <b>Purity/Specificity:</b> | Anti-FLIP Antibody is purified by DEAE purification methods. FLIP has short form (FLIPS) and long form (FLIPL) and antibody recognizes the FLIPL only.                        |
| <b>Relevant Links:</b>     | <ul style="list-style-type: none"><li>• <a href="#">UniProtKB - O35732</a></li><li>• <a href="#">GenelD - 12633</a></li><li>• <a href="#">NCBI - NP_001276633.1</a></li></ul> |

## Application Details

|                             |  |
|-----------------------------|--|
| <b>Tested Applications:</b> | ELISA, IHC, WB   |
| <b>Application Note:</b>    | Anti-FLIP Antibody has been tested for use in ELISA, Western Blotting, Immunocytochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 55 kDa in Western Blots of specific cell lysates and tissues. |
| <b>Assay Dilutions:</b>     | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.  |
| <b>ELISA:</b>               | 1:5,000  |
| <b>WB:</b>                  | 0.5 µg/mL  |

## Formulation


|                        |   |
|------------------------|---|
| <b>Physical State:</b> | Liquid (sterile filtered)                               |
| <b>Concentration:</b>  | 1 mg/mL by UV absorbance at 280 nm                      |
| <b>Buffer:</b>         | 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2 |
| <b>Preservative:</b>   | 0.02% (w/v) Sodium Azide                                |
| <b>Stabilizer:</b>     | None  |

## Shipping & Handling

|                            |   |
|----------------------------|---|
| <b>Shipping Condition:</b> | Dry Ice   |
| <b>Storage Condition:</b>  | Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |


**Expiration:** Expiration date is one (1) year from date of receipt.

## Images



### Immunohistochemistry

Immunocytochemistry of Rabbit anti-FLIP antibody. Cell Type: 3T3 cells. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: anti-FLIP antibody at 5 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: FLIP is cytoplasmic. Staining: FLIP as precipitated brown signal with hematoxylin purple nuclear counterstain.



### Western Blot

Western Blot of Rabbit anti-FLIP antibody. Lane 1: NIH/3T3 whole cell lysate. Load: 35 µg per lane. Primary antibody: anti-FLIP antibody at 1:500 for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 55.3 kDa, 55 kDa for FLIP. Other band(s): FLIP splice variants and isoforms.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.