

Datasheet for 200-401-FF4**TIRP Antibody****Overview**

Description:	Anti-TIRP (RABBIT) Antibody - 200-401-FF4
Item No.:	200-401-FF4
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	TIRP is a member of the Toll/interleukin-1 receptor (TIR) family, a group of proteins that include the Toll-like receptors (TLRs). TLRs are signaling molecules that recognize different pathogen-associated molecular patterns (PAMPs) and serve as an important link between the innate and adaptive immune responses. TIRP, along with other molecules such as TRIF, MAL, and MyD88, serves as an adaptor protein that allows for the interaction and activation of the IL-1R-associated kinase (IRAK) family, the subsequent activation of TNF receptor associated factor (TRAF)-6, and ultimately the activation of NF-κB. Expression of TIRP appears to be essential for TLR4 signaling.
Synonyms:	TIRP Antibody, TIRP, TRAM, TIRAP3, MyD88-4, TICAM-2, TIRP, TIR domain-containing adapter molecule 2, Putative NF-kappa-B-activating protein 502
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	TICAM2
Reactivity:	Human
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-TIRP antibody was prepared from whole rabbit serum produced by repeated immunizations with a peptide corresponding to 16 amino acids at the C-terminus of human TIRP.

Purity/Specificity:	Anti-TIRP Antibody is purified by ion exchange chromatography. Cross reactivity with TIRP from other sources has not been determined.
Relevant Links:	• UniProtKB - Q86XR7 • GeneID - 100302736 • NCBI - NP_001157940.1

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-TIRP Antibody has been tested for use in ELISA, Western Blotting, Immunocytochemistry and Immunofluorescence. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 27 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000
IF:	10 µg/mL
WB:	1 µg/mL


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling


Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images




Immunohistochemistry

Immunocytochemistry of Rabbit anti-TIRP antibody. Cell Type: Raji cells. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: TIRP antibody at 10 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: TIRP is located in the cellular membrane. Staining: TIRP is stained brown with hematoxylin purple nuclear counterstain.



Immunofluorescence Microscopy

Immunofluorescence Microscopy of TIRP antibody. Cell Type: Raji cells. Fixation: 0.5% PFA. Antigen retrieval: not required. Primary antibody: TIRP antibody at 10 µg/mL for 1 h at RT. Secondary antibody: Fluorescein rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: TIRP is located in the cell membrane. Staining: TIRP as green fluorescent signal.



Western Blot

Western Blot of TIRP antibody. PC-3 cell lysate. Load: 35 µg per lane. Primary antibody: TIRP antibody at 1 µg/ml for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 42 kDa, 27 kDa for TIRP. Other band(s): TIRP splice variants and isoforms.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.