

Datasheet for 200-401-FF7**TLR1 Antibody****Overview**

Description:	Anti-TLR1 (RABBIT) Antibody - 200-401-FF7
Item No.:	200-401-FF7
Size:	100 µg
Applications:	ELISA, IHC, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Toll-like receptors (TLRs) are evolutionarily conserved pattern-recognition molecules resembling the toll proteins that mediate antimicrobial responses in <i>Drosophila</i> . These proteins recognize different microbial products during infection and serve as an important link between the innate and adaptive immune responses. The TLRs act through adaptor molecules such as MyD88 and TIRAP to activate various kinases and transcription factors so the organism can respond to potential infection. TLR1 is co-expressed with TLR2 on myeloid cells of the innate immune systems in lymphoid tissue such as monocytes and dendritic cells where they form heterodimers that can recognize triacylated lipoproteins.
Synonyms:	TLR1 Antibody, TIL, CD281, rsc786, TIL. LPRS5, KIAA0012, Toll-like receptor 1, TIL
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	TLR1
Reactivity:	Human
Immunogen Type:	Conjugated Peptide
Immunogen:	Anti-TLR1 antibody was prepared from whole rabbit serum produced by repeated immunizations with a peptide corresponding to 17 amino acids near the internal region of human TLR1.

Purity/Specificity:	Anti-TLR1 Antibody is affinity chromatography purified via peptide column. Cross reactivity with TLR1 from other sources has not been determined.
Relevant Links:	• UniProtKB - Q15399 • GeneID - 7096 • NCBI - NP_003254.2

Application Details

Tested Applications:	ELISA, IHC, WB
Application Note:	Anti-TLR1 Antibody has been tested for use in ELISA, Western Blotting, Immunocytochemistry. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 90 kDa in Western Blots of specific cell lysates and tissues.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000-1:20,000
WB:	1-2 µg/mL


Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling


Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Immunohistochemistry

Immunocytochemistry of TLR1 antibody. Cell Type: Daudi cells. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: TLR1 antibody at 10 µg/mL for 1 h at RT. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Localization: TLR1 is located in the cytoplasm, endosome, lipid droplet, and cell membrane. Staining: TLR1 is stained brown with hematoxylin purple nuclear counterstain.



Western Blot

Western Blot of TLR1 antibody in Daudi cell lysate. Lane A: TLR1 antibody at 1 µg/mL. Lane B: TLR1 antibody at 2 µg/mL. Load: 35 µg per lane. Primary antibody: TLR1 antibody at designated concentrations for overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 47 kDa, 95 kDa for TLR1. Other band(s): TLR1 splice variants and isoforms.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.