

Datasheet for 200-401-H27**Trail Antibody****Overview**

Description:	Anti-TRAIL (RABBIT) Antibody - 200-401-H27
Item No.:	200-401-H27
Size:	100 µg
Applications:	ELISA, IF, IHC, WB
Reactivity:	Human
Host Species:	Rabbit

Product Details

Background:	Trail Antibody detects cytokines in the TNF family. Apoptosis, or programmed cell death, occurs during normal cellular differentiation and development of multicellular organisms. Apoptosis is induced by certain cytokines including TNF and Fas ligand in the TNF family through their death domain containing receptors, TNFR1 and Fas. A novel member in the TNF family was recently identified and designated TRAIL (for TNF-related apoptosis-inducing ligand) and Apo-2L (for Apo-2 ligand) ¹ , TRAIL is a type II membrane protein and expressed in a variety of human tissues. Two novel death domain containing receptors DR4 and DR5 have been identified as the receptor for TRAIL ³⁻⁶ . Like TNF and Fas ligand, TRAIL induces apoptosis and NF-κB activation in many tissues and cells. Anti-Trail antibodies are ideal for investigators involved in Cytokines and Growth Factors and Apoptosis research.
Synonyms:	Apo-2L, TRAIL
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	TNFSF10
Reactivity:	Human
Immunogen Type:	Conjugated Peptide

Immunogen:	Trail Antibody was produced from whole rabbit serum prepared by repeated immunizations with a peptide corresponding to amino acids near the c-terminus of human TRAIL.
Purity/Specificity:	Anti-Trail Antibody is affinity chromatography purified via peptide column. A BLAST analysis was used to suggest cross-reactivity with Trail with Human based on 100% homology with the immunizing sequence. Cross-reactivity with Trail from other sources has not been determined.
Relevant Links:	• NCBI - NP_003801 • UniProtKB - P50591 • GeneID - 8743

Application Details

Tested Applications:	ELISA, IF, IHC, WB
Application Note:	Anti-TRAIL Antibody is tested for use in E, WB, IF, and IHC. Expect a band approximately ~32.5 kDa on specific lysates. Western Blot, Immunohistochemistry, and Immunofluorescence tested in human samples. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
IHC:	20µg/mL
WB:	1 µg/mL

Formulation


Physical State:	Liquid (sterile filtered)
Concentration:	1 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.02% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Dry Ice
Storage Condition:	Store vial at -20° C prior to opening. Aliquot contents and freeze at -20° C or below for extended storage. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration:

Expiration date is one (1) year from date of receipt.


Images**Immunofluorescence Microscopy**

Immunofluorescence of TRAIL.

Tissue: human brain tissue.

Primary: Anti-Trail antibody at 20 µg/mL.


Staining: Trail Antibody (red), DAPI (blue).

**Immunohistochemistry**

Immunohistochemistry of TRAIL.

Tissue: human brain tissue.

Antibody: Anti-TRAIL antibody at 20 µg/mL.

**Western Blot**

Western blot analysis of TRAIL.

Lysate: HeLa cell lysate containing 10, 2.5, or 1 ng of recombinant protein containing extracellular domain of TRAIL.

Primary: Anti-TRAIL antibody at 1 µg/mL.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.