

## Datasheet for 200-4135

## Alkaline Phosphatase Antibody

### Overview

<b>Description:</b>	Anti-Alkaline Phosphatase (Calf Intestine) (RABBIT) Antibody (BULK ORDER) - 200-4135
<b>Item No.:</b>	200-4135
<b>Size:</b>	50 mg
<b>Applications:</b>	WB, IF, IHC, IP
<b>Reactivity:</b>	Bovine
<b>Host Species:</b>	Rabbit

### Product Details

<b>Background:</b>	In most mammals there are four different isozymes: placental, placental-like, intestinal and tissue non-specific (liver/bone/kidney). Alkaline Phosphatase catalytic activity is phosphate monoester + H <sub>2</sub> O = an alcohol + phosphate.
<b>Synonyms:</b>	rabbit anti-Alkaline Phosphatase Antibody, Alkaline phosphomonoesterase antibody, ALPI antibody, Glycerophosphatase antibody, IAP antibody, Intestinal alkaline phosphatase antibody, Kasahara isozyme antibody
<b>Host Species:</b>	Rabbit
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	IgG

### Target Details

<b>Gene Name:</b>	ALP1
<b>Reactivity:</b>	Bovine
<b>Immunogen Type:</b>	Native Protein
<b>Immunogen:</b>	Alkaline Phosphatase [Calf Intestine]

**Purity/Specificity:** Anti-Alkaline Phosphatase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as purified and partially purified Alkaline Phosphatase [Calf Intestine]. Cross reactivity against Alkaline Phosphatase from other tissues and species may occur but have not been specifically determined.

- Relevant Links:**
- [200-4135 SDS](#)
  - [UniProtKB - P19111](#)
  - [NCBI - P19111.2](#)
  - [GenelD - 280993](#)

## Application Details

<b>Tested Applications:</b>	WB
<b>Suggested Applications:</b>	IF, IHC, IP (Based on references)
<b>Application Note:</b>	Anti-Alkaline Phosphatase has been tested against 1.0 ug of Alkaline Phosphatase [Calf intestine] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:2,000 to 1:8,000 of the reconstitution concentration is suggested for this product.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
<b>ELISA:</b>	1:20,000 - 1:100,000
<b>IHC:</b>	1:1,000 - 1:5,000
<b>WB:</b>	1:2,000 - 1:10,000

## Formulation

<b>Physical State:</b>	Lyophilized
<b>Concentration:</b>	10.0 mg/mL by UV absorbance at 280 nm
<b>Buffer:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
<b>Preservative:</b>	0.01% (w/v) Sodium Azide
<b>Stabilizer:</b>	None
<b>Reconstitution Volume:</b>	5.0 mL

**Reconstitution Buffer:** Restore with deionized water (or equivalent)


## Shipping & Handling

**Shipping Condition:** Ambient

**Storage Condition:** Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Expiration:** Expiration date is one (1) year from date of receipt.

## Images



### Western Blot

Western blotting of secretory alkaline phosphatase (SEAP) expressed in INS-1  $\beta$ -cells. (A) A stably transfected clone of INS-1 cells (clone 12, Trnfxn) immunoblotted for alkaline phosphatase next to untransfected control cells (Con). In addition, a faster-migrating specific band is detected in the stably transfected cells. Upon glycan digestion (+), the steady state distribution of SEAP is divided into endo-H-sensitive (S) and endo-H-resistant (R) forms, the latter indicative of a significant Golgi and/or post-Golgi pool of the protein in these cells. (B) A comparison of SEAP expression level in the clone shown in panel A (C-12) to two other independently selected INS-1 clones (C-14 and C-15), each of which maintains a similar proportion of endo-H-resistant SEAP. Fig 1.

PMID: 16608874



### Western Blot



Acquisition of endo H resistance and secretion of SEAP at 2 hours of chase in INS-1  $\beta$ -cells. Cells were metabolically labeled for 30 minutes and the media (M), cells (C) or total (T, media + cells) were immunoprecipitated with anti-alkaline-phosphatase antibody. At the zero chase time, SEAP had not yet appeared in the medium and all intracellular SEAP was sensitive to digestion with endo H (S). At 2 hours of chase, more than half of labeled SEAP had acquired endo H resistance (R); 35% of labeled SEAP reached the medium (all of which is endo H resistant), and 25% of intracellular SEAP accumulated as an endo-H-resistant form. The data are representative of three experiments. Fig 2.

PMID: 16608874

### Immunofluorescence Microscopy


Preservation of enterocytes and enteroendocrine cells and cell polarity in  $G\alpha q/11$ -deficient mice. Immunohistochemical studies were performed for intestinal alkaline phosphatase (enterocyte marker), chromogranin A (endocrine cell marker), F-actin (mainly localized in the subapical membrane), and E-cadherin (localized in the basolateral membrane). No differences were observed between control and Int-Gq/G11 DKO mice. Scale bars: 200  $\mu$ m. Bottom: Analysis of the number of chromogranin A-positive cells (arrowheads) in crypt and villi in the 4 genotypes. Cell numbers are graphed as the means  $\pm$  SD (30 crypts and villi/mouse, n = 4 per genotype). Fig 4.

PMID: 28174748



### Western Blot

Western Blot of rabbit anti Alkaline Phosphatase antibody  
Lane 1: Calf intestine Alkaline Phosphatase. Load:  $\sim$ 1.5  $\mu$ g  
Block: 5% Blotto 1 hour at 4°C. Primary antibody: diluted to 1:1000 in 5% Blotto overnight at 4°C. Secondary antibody: Goat anti Rabbit 611-103-122 Lot#21231 1:20,000 in MB-070 1 Hour 4°C Substrate: Femtomax 110.

**Western Blot**

Western Blot of Rabbit anti-Alkaline Phosphatase (Calf Intestine) Antibody. Lane 1: Alkaline Phosphatase (Calf Intestine). Load: 50 ng per lane. Primary antibody: Rabbit anti-Alkaline Phosphatase (Calf Intestine) Antibody at 1:1,000 overnight at 4°C. Secondary antibody: HRP rabbit secondary antibody (p/n 611-103-122) at 1:40,000 for 30 min at RT. Block: (p/n MB-070) for 30 min at RT.

**References**

- Iwaki A. et al. Loss of Rab6a in the small intestine causes lipid accumulation and epithelial cell death from lactation. *FASEB J.* (2020)
- Jones et al. Cellular Plasticity of Defa4Cre-Expressing Paneth Cells in Response to Notch Activation and Intestinal Injury. *Cellular and Molecular Gastroenterology and Hepatology* (2018)
- Teoh JJ et al. BIG1 is required for the survival of deep layer neurons, neuronal polarity, and the formation of axonal tracts between the thalamus and neocortex in developing brain. *PLoS One.* (2017)
- Watanabe N et al. Requirement of Gαq/Gα11 signaling in the preservation of mouse intestinal epithelial homeostasis. *Cell Mol Gastroenterol Hepatol.* (2016)
- Sobajima, T et al. Rab11a is required for apical protein localisation in the intestine. *Biology Open* (2014)
- Paladino S et al. Golgi sorting regulates organization and activity of GPI proteins at apical membranes. *Nat Chem Biol.* (2014)
- Lara-Lemus R et al. Luminal protein sorting to the constitutive secretory pathway of a regulated secretory cell. *J Cell Sci.* (2006)

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.