

Datasheet for 200-4137

Glucose Oxidase Antibody

Overview

Description:	Anti-Glucose Oxidase (RABBIT) Antibody (BULK ORDER) - 200-4137
Item No.:	200-4137
Size:	50 mg
Applications:	ELISA, WB
Reactivity:	Aspergillus niger
Host Species:	Rabbit

Product Details

Background:	Anti-Glucose Oxidase detects glucose oxidase. Glucose oxidase enzyme (GOx) is an oxido-reductase that catalyzes the oxidation of glucose to hydrogen peroxide and D-glucono- δ -lactone. In cells, it aids in breaking the sugar down into its metabolites. Glucose oxidase is widely used for the determination of free glucose in body fluids, in vegetal raw material, and in the food industry. It also has many applications in biotechnologies, typically enzyme assays for biochemistry including biosensors in nanotechnologies. Anti-Glucose Oxidase Antibody is ideal for investigators involved in Cell Signaling and Signal Transduction research.
Synonyms:	rabbit anti-Glucose Oxidase Antibody, Beta D Glucose Oxygen 1 Oxido Reductase antibody, Glucose oxidase (Precursor) antibody, Glucose Oxyhydrase antibody, GOD antibody
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	gox
Reactivity:	Aspergillus niger
Immunogen Type:	Native Protein
Immunogen:	Anti-Glucose Oxidase Antibody was produced by repeated immunizations with Aspergillus niger Glucose Oxidase protein.

Purity/Specificity: Anti-Glucose Oxidase is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as purified and partially purified Glucose Oxidase [*Aspergillus niger*]. Cross reactivity against Glucose Oxidase from other tissues and species may occur but have not been specifically determined.

Relevant Links:

- [UniProtKB - P13006](#)

Application Details

Tested Applications: ELISA, WB

Application Note: Anti-Glucose Oxidase Antibody has been tested by ELISA and western blot and is suitable for IHC. Researchers should determine optimal titers for applications that are not stated below.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

ELISA: 1:35,000 - 1:160,000

IHC: 1:2,000 - 1:8,000

WB: 1:3,500 - 1:16,000

Formulation

Physical State: Lyophilized

Concentration: 10.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: 0.01% (w/v) Sodium Azide

Stabilizer: None

Reconstitution Volume: 5.0 mL

Reconstitution Buffer: Restore with deionized water (or equivalent)


Shipping & Handling

Shipping Condition: Ambient

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Rabbit anti-Glucose Oxidase Antibody.
Lane 1: Glucose Oxidase. Load: 50 ng per lane. Primary antibody: Glucose Oxidase antibody at 1:1,000 for overnight at 4°C. Secondary antibody: DyLight™ 649 rabbit secondary antibody (p/n 611-143-002) at 1:20,000 for 30 min at RT. Block: (p/n MB-070) for 30 min at RT. Predicted/Observed size: 80 kDa, 80 kDa for Glucose Oxidase from *Aspergillus niger*. Other band(s): None.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.