

Datasheet for 200-4182**Urease Antibody****Overview**

Description:	Anti-Urease (Jack Bean) (RABBIT) Antibody (BULK ORDER) - 200-4182
Item No.:	200-4182
Size:	50 mg
Applications:	WB
Reactivity:	Jack Bean
Host Species:	Rabbit

Product Details

Background:	Urea hydrolase involved in nitrogen recycling from ureide, purine, and arginine catabolism. Urease catalyzes the hydrolysis of urea to produce ammonia and carbamate (ammonia and carbonic acid). Ureases are found in numerous bacteria, fungi, algae, invertebrates, plants, in soils, as a soil enzyme. Due to the catalytic activity producing ammonia the pH of the environment increases. Ureases, when bacterial, can be a mode for the pathogens of many medical conditions such as hepatic encephalopathy, hepatic coma, urinary stones, and peptic ulcers. Urea is found naturally in the environment as well as artificially introduced. Anti-Urease antibody is ideal for researchers interested in ecosystems, microbes, and gut bacteria.
Synonyms:	rabbit anti-Urease Antibody, Urea Amidohydrolase antibody
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	UREA_CANEN
Reactivity:	Jack Bean
Immunogen Type:	Native Protein
Immunogen:	Urease [Jack Bean]

Purity/Specificity: Anti-Urease is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as purified and partially purified Urease [Jack Bean]. Cross reactivity against Urease from other tissues and species may occur but have not been specifically determined.

Relevant Links:

- [UniProtKB - P07374](#)
- [NCBI - AAA83831.1](#)

Application Details

Tested Applications:	WB
Application Note:	Anti-Urease Antibody has been tested in western blot and suitable against 1.0 ug of Urease [Jack Bean] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Rabbit IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:15,000 to 1:60,000 of the reconstitution concentration is suggested for this product.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:5,000 - 1:20,000
IP:	1:100
WB:	1:500 - 1:5,000

Formulation


Physical State:	Lyophilized
Concentration:	10.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None
Reconstitution Volume:	5.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition: Ambient

Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Rabbit anti-Urease (Jack Bean) Antibody.
 Lane 1: Urease (Jack Bean). Load: 50 ng per lane. Primary antibody: Urease primary antibody at 1:1,000 overnight at 4°C. Secondary antibody: Peroxidase rabbit secondary antibody (p/n 611-103-122) at 1:40,000 for 30 min at RT. Block: (p/n MB-070) for 30 min at RT. Predicted/Observed size: 90 kDa, 90 kDa for Urease.

References

- Ramlau R et al. Safety of unconventional antibody-drug conjugate L-DOS47 in a phase I/II monotherapy study targeting advanced NSCLC. *Front Oncol.* (2025)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.