

Datasheet for 200-4195

Streptavidin Antibody

Overview

Description:	Anti-Streptavidin (RABBIT) Antibody (BULK ORDER) - 200-4195
Item No.:	200-4195
Size:	50 mg
Applications:	Dot Blot, ELISA, WB
Reactivity:	Streptavidin
Host Species:	Rabbit

Product Details

Background:	Streptavidin antibody is a 60 kDa protein purified from the bacterium <i>Streptomyces avidinii</i> . Streptavidin homo-tetramers have an extraordinarily high affinity for biotin (also known as vitamin B7). With a dissociation constant (Kd) on the order of $\approx 10^{-14}$ mol/L, [1] the binding of biotin to streptavidin is one of the strongest non-covalent interactions known in nature. Streptavidin is used extensively in molecular biology and bio-nanotechnology due to the streptavidin-biotin complex's resistance to organic solvents, denaturants (e.g. guanidinium chloride), detergents (e.g. SDS, Triton), proteolytic enzymes, and extremes of temperature and pH. Anti-STREPTAVIDIN antibody is ideal for investigators involved in GFP and Epitope research.
Synonyms:	rabbit anti-Streptavidin Antibody, rabbit anti Streptavidin
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Streptavidin
Immunogen Type:	Native Protein
Immunogen:	Streptavidin [<i>Streptomyces avidinii</i>]

Purity/Specificity:	Anti-Streptavidin antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum as well as purified and partially purified Streptavidin [<i>Streptomyces avidinii</i>]. No cross reactivity occurs against Avidin.
----------------------------	---

Application Details

Tested Applications:	Dot Blot, ELISA, WB
Application Note:	Anti-Streptavidin antibody has been tested by ELISA, dot blot, and western blot, and is suitable for immunohistochemistry, as well as other assays requiring lot-to-lot consistency.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:200,000
IHC:	1:1,500 - 1:5,000
WB:	1:3,000 - 1:12,000

Formulation


Physical State:	Lyophilized
Concentration:	10.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None
Reconstitution Volume:	5.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Rabbit Anti-Streptavidin Antibody.

Lane 1: Streptavidin - Reduced (p/n S000-01) [0.05µg].

Lane 2: Opal Prestained Molecular Weight Marker (p/n MB-210-0500).


Lane 3: Streptavidin - Non-Reduced (p/n S000-01) [0.05µg].

Primary Antibody: Rabbit Anti-Streptavidin at 1:1000 overnight at 2-8°C.

Secondary Antibody: Goat Anti-Rabbit IgG Peroxidase (p/n 611-103-122) at 1:70,000 for 30 mins at RT.

Block: Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) for 60 mins at RT.

Expect: ~13kDa reduced, ~60kDa non-reduced.



Immunofluorescence Microscopy

CD8 staining efficiency. DTH biopsy from patient 5 (lesion DC + 3 peptides) stained with HLA-A2.1-

tyrosinase369 tetramer (red), anti-CD8 (green) and DAPI (blue) with increasing exposure time for the FITC filter

(ranging from 2–3 s (a) to 15–20 s (c), for comparison: exposure time TRITC 7–8 s). Arrows indicate CD8+ cells that

were not visible at shorter exposure times for the FITC filter.

In c, only for the boxed area an increased exposure time

(15–20 s) is shown to prevent overexposure from other parts

of the section. d Enlarged two color overlay for green (CD8)

and blue (DAPI) of the boxed area from c. Tetramer analysis

by flow cytometry of DTH-derived T cells from patient 5. On

the y-axis MHC class I tetramer PE staining and on the x-axis

CD4 (e) or CD8 (f). Note the difference in fluorescence

intensity in the tetramer positive population (MFI CD4:152

vs. MFI CD8:83) whereas the percentage of positive cells is

equal (17%). CD8+ cells visualized with the ABC-peroxidase


kit with AEC (red-brown staining) as substrate in

corresponding areas on tissue sections from tonsil fixed with

acetone (g) or PFA (h).

Fig 5

PMID: 17440724





Western Blot

Rockland Rabbit anti Streptavidin (200-4195 lot 23495) and Biotin conjugated Rabbit anti-trypsin inhibitor antibody (200-4679 lot 6594) were used to detect target proteins Trypsin Inhibitor (left) and Streptavidin (right) under reducing (R) and non-reducing (NR) conditions. Reduced samples of purified target proteins contained 4% BME and were boiled for 5 minutes. Samples of ~1ug of protein per lane were run by SDS-PAGE. Protein was transferred to nitrocellulose and probed with 1:1000 dilution of primary antibody (ON 4 C). Detection shown was using Dylight 649 conjugated Donkey anti rabbit (611-743-127 lot 20831 1:10K 1.5 hr RT in MB-070) and imaged on the BioRad VersaDoc System.

Dot Blot

Dot Blot showing the detection of Streptavidin. A three-fold serial dilution of DyLight™ 488 Conjugated Streptavidin starting at 200ng was spotted onto 0.45 µm nitrocellulose and blocked in 1% BSA-TTBS (p/n MB-013, diluted to 1X) 30 min at 20°C. Anti-Streptavidin (RABBIT) Antibody (p/n 200-401-095) was incubated in Blocking Buffer for Fluorescent Western Blotting (p/n MB-070) at 1:1,000 for 1 Hour at 20°C. An HRP Gt-a-Rb secondary antibody (p/n 611-103-122) was incubated at 1:40,000 for 30 min at 20°C and imaged using the Bio-Rad VersaDoc® 4000 MP.



Western Blot

Western Blot results of Rabbit Anti-Streptavidin Antibody. Lane 1: Opal Prestained Molecular Weight Ladder (p/n MB-210-0500). Lane 2: Streptavidin (p/n S000-01). Load: 50ng. Primary Antibody: Rabbit Anti-Streptavidin at 1:1000 overnight at 4°C. Secondary Antibody: Goat anti-Rabbit HRP (p/n 611-103-122) at 1:70,000 for 30min at RT. Blocking: BlockOut (p/n MB-073) for 30 min at RT.

References

- Koyroytsaltis-McQuire DJP et al. Metasurface analogues of molecular diastereomers from hierarchical multiscale chiral interactions with biomolecules. *Nat Commun.* (2026)
- M Herrmann et al. Enzymatically-generated fluorescent detection in micro-channels with internal magnetic mixing for the development of parallel microfluidic ELISA. *Lab Chip.* (2006)
- Shiyankovskii SV et al. Real-time microbe detection based on director distortions around growing immune complexes in lyotropic chromonic liquid crystals. *Phys Rev E Stat Nonlin Soft Matter Phys.* (2005)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.